



Original

Association between body mass index and cardiorespiratory fitness as predictor of health status in schoolchildren



S.S. De Araujo^{a,*}, R. Miguel-dos-Santos^{a,b}, R.J.S. Silva^{a,b}, A.C. Cabral-de-Oliveira^{a,b}

^a Physiology Department, Federal University of Sergipe, Brazil

^b Nucleus Postgraduate in Physical Education, Federal University of Sergipe, Brazil

ARTICLE INFO

Article history:

Received 8 August 2013

Accepted 18 February 2014

Keywords:

Cardiorespiratory fitness

Body composition

Body mass index

Fitnessgram[®]

Schoolchildren

ABSTRACT

Objective: Evaluate the cardiorespiratory fitness (CRF) and the body mass index (BMI) of schoolchildren according to the cutoff points established by the Fitnessgram[®]. Just as determine the influence of BMI on CRF in children and propose an equation for predicting VO₂max from the body composition.

Method: The sample consisted of 288 students of basic level of public education in the city of Aracaju – Sergipe – Brazil, aged between 10 and 14 years, of these, 142 were female. To evaluate the CRF the 20-m shuttle run test was chosen. The results of the variable were classified in accordance with the criteria established by the Fitnessgram[®].

Results: It was evidenced that the cardiorespiratory fitness on males was higher than females. Relative to BMI, there were no differences between the groups, while more than 80% of the entire sample behaved within criteria established by the Fitnessgram[®]. On the other hand, 57% of the female group presented VO₂max below criterion. It was confirmed an inverse relationship between VO₂max and BMI on children and adolescents.

Conclusion: The students in this study showed adequate levels of body composition and CRF below the area related to health, according to the criteria of the Fitnessgram[®]. The proposition of a linear regression equation to estimate VO₂max without exercise pointed out an inverse relationship between both variables.

© 2013 Consejería de Educación, Cultura y Deporte de la Junta de Andalucía. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Asociación entre el índice de masa corporal y la capacidad cardiorrespiratoria como predictor del estado de salud de escolares

RESUMEN

Objetivo: Evaluar la aptitud cardiorrespiratoria y el índice de masa corporal (IMC) de escolares mediante los puntos de corte establecidos por la batería internacional de test Fitnessgram[®]. Así como determinar la influencia del IMC en la aptitud cardiorrespiratoria (CRF) en niños y proponer una ecuación de predicción del VO₂máx a partir de la composición corporal.

Método: La muestra estuvo conformada por 288 estudiantes de nivel básico de la educación pública de la ciudad de Aracaju-Sergipe-Brasil, de edades comprendidas entre 10-14 años, de los cuales, 142 eran mujeres. Para evaluar el CRF fue utilizada la prueba de 20-m shuttle run test (PCN). Los resultados de la variable se clasificaron de acuerdo con los criterios de aplicación establecidos por el test internacional Fitnessgram[®].

Resultados: Se evidenció que la CRF en varones fue mayor que en las mujeres. Relativo al IMC, no hubo diferencias entre los grupos, mientras que más de 80% de toda la muestra se comportó dentro de los criterios establecidos por la Fitnessgram[®]. Por otro lado, el 57% del grupo femenino presentó VO₂máx por debajo de los niveles. Se confirmó una relación inversa entre el VO₂máx y el IMC en niños y adolescentes.

Conclusión: Los escolares, en este estudio, mostraron niveles adecuados de composición corporal y CRF por debajo del área relacionada con la salud de acuerdo con los criterios de Fitnessgram[®]. La proposición de

Palabras clave:

Aptitud cardiorrespiratoria

Composición corporal

Índice de masa corporal

Fitnessgram[®]

Escolares

* Corresponding author.

E-mail address: silvan.ssa@gmail.com (S.S. De Araujo).

una ecuación de regresión lineal, para estimar el VO₂máx sin ejercicio señaló una relación inversa entre ambas variables.

© 2013 Consejería de Educación, Cultura y Deporte de la Junta de Andalucía. Publicado por Elsevier España, S.L.U. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Associação entre índice de massa corporal e aptidão cardiorrespiratória como preditor de status de saúde em escolares

R E S U M O

Palavras-chave:

Aptidão cardiorrespiratória
Composição corporal
Índice de massa corporal
Fitnessgram®
Escolares

Objetivo: Avaliar a aptidão cardiorrespiratória e índice de massa corporal (IMC) de escolares de acordo com os pontos de corte estabelecidos pela bateria de teste internacional Fitnessgram®. Além de determinar a influência do IMC sobre a aptidão cardiorrespiratória em crianças e propor uma equação de predição do VO₂máx a partir da composição corporal.

Método: A amostra consistiu de 288 estudantes do ensino básico da educação pública na cidade de Aracaju (Sergipe, Brasil), com idade entre 10-14 anos, dos quais 142 eram meninas. Para avaliar a aptidão cardiorrespiratória optou-se pelo 20-m shuttle run test. Os resultados das variáveis foram classificados de acordo com os critérios estabelecidos pelo Fitnessgram®.

Resultados: Constatou-se que a aptidão cardiorrespiratória dos meninos foi maior do que das meninas. Com relação ao IMC, não houve diferenças entre os grupos, enquanto mais de 80% de toda a amostra se comportou dentro dos critérios estabelecidos pelo Fitnessgram®. Por outro lado, 57% do grupo feminino apresentou VO₂máx abaixo dos critérios. Confirmou-se uma relação inversa entre VO₂máx e o IMC em crianças e adolescentes.

Conclusão: Os alunos deste estudo apresentaram níveis de composição corporal adequados e de aptidão aeróbia abaixo da área relacionada à saúde, de acordo com os critérios do Fitnessgram®. A proposição de uma equação de regressão linear para estimar o VO₂máx sem exercício apontou uma relação inversa entre as 2 variáveis.

© 2013 Consejería de Educación, Cultura y Deporte de la Junta de Andalucía. Publicado por Elsevier España, S.L.U. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

The decrease in the amount of motor activity observed on children and adolescents in different socioeconomic spectra¹ has raised interest in the knowledge of physical fitness and health, in order to classify it from scores based on referenced criteria both nationally and internationally, which involve physical qualities able to reflect patterns of behavior and environmental.

Closely related to the health, the low levels of cardiorespiratory fitness (CFR) on childhood and adolescence are associated with an increased early death risk due to any cause, especially cardiovascular diseases.^{2,3} So the prevention of cardiovascular diseases should begin still during the school stage, given that the adoption of healthy attitudes on that age range is perpetuated into adulthood.

In this way it is important to detect in advance, even within the school environment, children and adolescents with a predisposition to low aerobic fitness. In this sense, the development of mathematical models for predicting CRF without exercise test could be feasible in epidemiological studies⁴ and very useful for physical education teachers.

From this perspective, the presented study aimed to evaluate the CRF and the BMI of schoolchildren of both sexes according to the cutoff points established by the Fitnessgram®, as well as determine the influence of BMI on CRF of children and propose an equation for predicting VO₂max from the body composition.

Method

Participants

Data collection was performed in a cross-sectional mode, and the target population of this study consisted on students of both genders enrolled in eight elementary schools distributed in the urban area of Aracaju, a city located on the coast of the state of

Sergipe, Brazil. It lies between the capitals with lower socioeconomic inequalities and contributes greatly to the second highest Human Developing Index (HDI) among the states in northeastern region. The intervention took place between the months of September and November 2005, in obedience to the school calendar of the season.

To determine the surveyed schools, the city was divided into three regions: north, center and south, and by means of a draw three schools were chosen by region. Schools should have the minimum physical structure for the tests (e.g. multi-sports court). The sample, selected randomly during physical education classes, composed by 288 children (142 females) with age range between 9.50 and 14.49 years. We used the decimal age children as Ross and Marfell-Jones,⁵ which accurately represents the interval between the date of birth and date of the test.

The present study adopted the following inclusion criteria: belong to the preset age group, attend school regularly and take part in physical education classes on the day and time scheduled. The exclusion criteria were: experience any clinical symptoms that prevented temporarily or permanently be subjected to the tests and do not agree with the terms of consent by the responsible person.

Two professionals and four physical education academics carried out tests and measurements. To avoid discrepancies between measurements, each team member performed the same activity during the intervention. Before data collection, a training program was conducted in the test battery.

Procedures and measures

Anthropometric measurements

Anthropometric variables whose values are presented in Table 1, were body mass (kg) and height (m), as well as measures of skinfolds, in conformity with established standards.⁵ The body fat percentage (BF%) was determined as proposed Lohman.⁶ We used

Table 1

Descriptive profile of the sample (mean and standard deviation) for the anthropometry and cardiorespiratory fitness.

Variables	All group (n = 288)	Male (n = 142)	Female (n = 146)
Age (years)	12.2 (1.1)	12.2 (1.2)	12.1 (1.1)
Height (cm)	149.6 (10.2)	149.2 (11.1)	150.0 (9.2)
Body mass (kg)	41.6 (10.3)	41.5 (10.7)	41.8 (9.9)
BMI (kg/m ²)	18.4 (3.0)	18.3 (2.8)	18.4 (3.2)
BF%	17.1 (8.3)	14.4 (7.9)	19.9 (7.7)*
VO ₂ max	38.0 (3.7)	38.8 (3.8)	37.1 (3.4)*

BMI: body mass index; BF%, body fat percentage; VO₂max: maximal oxygen uptake.
* $p < 0.05$, statistically significant difference between male and female.

the caliper skinfold Cescorf[®] (CSC). The CSC presents identical design and mobility to the Harpenden model, provides a pressure of approximately 10 g/mm², measures to an accuracy of 0.1 mm, up to a jaw width of 88 mm.⁷ Skinfolts considered for the study were: triceps and subscapularis.

The body mass index (BMI) was calculated as the ratio of the body mass (kg) and the squared height (m²). Therefore, the cutoff points of the Fitnessgram[®] were used to calculate the prevalence of children and adolescents located in the recommended range and in the range of risk of overweight and obesity.

Cardiorespiratory fitness (CRF)

To evaluate the CRF from the maximal oxygen uptake (VO₂max), we opted for the 20-m shuttle run test, according to what Léger et al.⁸ determine. This test was chosen because of its affordable implementation and compliance with the motor tests related to health contained in the Fitnessgram[®] battery, created by the Cooper Institute for Aerobics Research⁹ and because of its reliability and validity on children and adolescents has been widely documented.^{8,10,11} VO₂max was calculated by the following equation, that covers the age range of this study. An r value of 0.71 was obtained:

$$y = 31.025 + 3.238x_1 - 3.248x_2 + 0.1536x_1x_2, \text{ where:}$$

$$y = \text{VO}_2\text{max (ml/kg/min)}$$

$$x_1 = \text{maximal speed attained (km/h)}$$

$$x_2 = \text{age (y)}$$

Fitnessgram[®] criteria

To classify the performance level regarding the criteria established by the Fitnessgram[®] (VO₂max: male 37.4–42.4, female 37.4–39.3; BMI: male 14.5–23.0, female 14.3–23.6),⁹ the results were categorized in three performance levels, “below”, “normal” and “above”. The following cutoff points were posted by Cooper Institute for Aerobics Research in 2013.

Ethics in research

This study was conducted after the approval of the Municipal Education and according to the laws in force and in compliance with the Guidelines and Standards for Research Involving Humans, it was approved by the Ethics Committee of the Federal University of Sergipe under registration 1138.0.000.107-06.

Statistical analysis

Descriptive statistics was used as first step to data analysis. A t -test for independent samples was chosen to compare means and the Chi-squared test was used to compare proportions. The relationship between VO₂max and BMI was verified using the Pearson linear correlation and the lineal regression analysis. Level of significance was set at $p < 0.05$. The results were analysed by using GraphPadPrism 5 or SPSS software when necessary.




Fig. 1. Relative values by sex, following the Fitnessgram cutoffs established for the Body Mass Index, BMI (kg/m²).

Results

The results of anthropometric variables observed in Table 1 were similar when comparing both sexes, regarding age, height, body mass (BM) and BMI. In terms of percentage of body fat, females had higher values ($t = 6.049$, $p < 0.0001$), and the boys showed higher values for maximum oxygen consumption ($t = -4.106$, $p < 0.0001$).

Regarding the behavior of BMI in both sexes (Fig. 1), it may be noted no differences based on established criteria. Considering the cutoff points for the Fitnessgram[®], 9.2% of girls were classified as “below”, 80.3% “normal,” and 10.5% “above”, the values considered in the same way for males were respectively, 8.2%, 79.5% and 12.3%.

In relation to VO₂max (Fig. 2), the percentage of boys classified as ‘above’, according to the cutoffs of the Fitnessgram[®], was higher (41.8%) than the percentage of girls (19.7%). No statistically significant difference was found between girls and boys in the other




Fig. 2. Relative values by sex, following the Fitnessgram cutoffs established for the VO₂max (ml kg⁻¹ min⁻¹). * $p < 0.05$: statistically significant difference between male and female; %: percentage; VO₂max: maximal oxygen uptake.




Fig. 3. Simple linear regression curve for the relationship between BMI and VO_2 max in males ($R^2 = 0.850$). VO_2 max: maximal oxygen uptake; BMI: body mass index.




Fig. 4. Simple linear regression curve for the relationship between BMI and VO_2 max in females ($R^2 = 0.907$). VO_2 max: maximal oxygen uptake; BMI: body mass index.

classifications. The percentage of boys and girls classified as 'below' were 44.5% and 57.1%, respectively. The percentage of boys and girls classified as 'normal' were 13.7% and 23.2%, respectively.

Both **Figs. 3 and 4** show the correlation between the two variables, as well as the influence of the independent variable, BMI exerts on the dependent, VO_2 max, by determining the simple linear regression and the R^2 value. This behavior was evaluated in both sexes and as many girls ($R^2 = 0.907$) as boys ($R^2 = 0.850$) presented strong correlation.

Analyzing **Table 2**, it was found in the study group a strong relationship in both sexes between the dependent and independent variables.

Table 2
Simple linear regression analysis to determine VO_2 max based on BMI.

	Pearson	R^2	<i>p</i>
Male			
VO_2 max = $-2.59(\text{BMI}) + 86.57$	-0.922	0.850	0.026
Female			
VO_2 max = $-0.39(\text{BMI}) + 32.86$	-0.953	0.907	0.012

Linear regression; VO_2 max: maximal oxygen uptake; BMI: body mass index.

Discussion

The present study aimed to evaluate the cardiorespiratory fitness and the BMI of schoolchildren of both sexes according to the cutoff points established by the Fitnessgram[®], as well as determine the influence of BMI on CRF of children and propose an equation for predicting VO_2 max from the body composition.

There was a low compliance to the Fitnessgram[®] criteria associated with risk of obesity (**Fig. 1**). These results were better than those found by Bergmann et al.,¹² in a study with students of 13–14 years old in the city of Canoas (Rio Grande do Sul, Brazil). In that occasion the authors detected 19% of boys and 15% of girls outside the health criteria recommended by Projeto Esporte Brasil (PROESP-BR). This may be attributed to the changes and differences in lifestyles of each region or country.¹³

In relation to other national studies, such as Hobold and Hübner¹⁴ with students of 10–14 years old, the BMI behaved as 'below', for both female and male from the city of Marechal Cândido Rondon (Paraná, Brazil). Ronque et al.¹⁵ in adolescents aged 10–15 years old from the city of Londrina (Paraná, Brazil), obtained higher BMI values than the present study, in individuals with low and high cardiorespiratory fitness based on the criteria of the Fitnessgram[®]. However, the sample here studied remained above the schoolchildren from the Vale do Cotinguiba (Sergipe, Brazil) studied by Petroski et al.¹⁶ Although with respect to Spanish 7–12 years old school children from Aragón, have shown to be equivalent.¹⁷

The investigation of morphological parameters becomes more relevant when detects possibilities of body fat accumulation, the emergence of chronic diseases and cardiovascular diseases. Malina and Bouchard¹⁸ emphasized that the female acquires greater gains of fat mass than boys from late childhood to late adolescence.

In this sense, the BF% (Table 1) of both groups were higher than those obtained by Dellagrana et al.¹⁹ and Hobold and Hübner.¹⁴ In this perspective, Lloyd et al.²⁰ in students aged 10–12 years old from Texas – USA, found a BF% of 14.3 in males, at the same level of this study, although the female showed 8.5%, a value very below the results of this study. Despite variations, these results put the sample of this study far from risk of overweight and obesity according Fitnessgram[®] criteria.²¹ This evidence was corroborated by the parameters of the BMI (**Fig. 1**), which also report levels below normal.

Several studies^{22–25} over the past decade have reported the trend towards a minor cardiorespiratory fitness on female school students compared to their male counterparts. This phenomenon can be explained by sudden changes in body composition during puberty, particularly fat mass for girls.¹⁸ In addition to this, LaMonte et al.²⁶ obtained that the maximal oxygen consumption is inversely proportional to BMI, in addition to other serum lipid parameters.

It was evidenced that the CRF of males was higher than females, while for the BMI, there were no differences between the groups. While most of the group remained in the normal range for BMI (79.9%), it drew attention to the high prevalence of the entire sample (50.8%) with poor performance for VO_2 max, particularly with respect to the female group, which 57.1% of them did not fit the health criteria adopted by Fitnessgram[®]. Furthermore, we proved the hypothesis of an inverse relationship between VO_2 max and BMI, in children and adolescents, a crucial period of life to develop these physical qualities.²⁷

Because it is an important functional aspect, which if neglected in childhood and adolescence leads to extreme damage to health in adulthood, the result of this study on CRF requires specific attention by the government and educational authorities. It is well known the secular trend in the decline of this physical quality by several factors, including increased time spent in sedentary behaviors, like a high rate of exposure to television.²⁸

This transition to sedentary lifestyle and its effects are strengthened by the phenomenon of urbanization of the society, as reported Chillón et al.,²⁹ who studied Spanish schoolchildren from antagonistic geographic areas, rural and urban, where lower levels of VO₂max were found in the urban school children compared to their rural peers. Data from two cross-sectional studies in adolescents of 12.5–17.5 years old from Zaragoza (Spain) revealed an increase in CRF over 5 years.³⁰ According to the authors, these data follow the opposite to the consistent decrease in cardiorespiratory fitness identified by studies around 27 countries and five geographic regions around the world.

The fact that the data from this study are aligned with the global trend is worrying, because low levels of this variable are related to risk of cardiovascular diseases, which begin in childhood and progress silently until adulthood, and are associated with a high number of deaths in some populations.^{22,31,32} The low performance of students from Aracaju (present study) on 20 m shuttle-run test – only one boy for every ten (13.7%) and one girl for every five (23.2%) met the cutoff points proposed by Fitnessgram® – contrasts with the results obtained by Guedes et al.,³³ who found that about 50% on both boys and girls, reached health criteria. According to the authors, these results are not optimistic, especially because they observed a decline in proportion when increasing age, especially in physical tests that involve large muscle groups.

Usually, girls tend to be more affected by the advancement of puberty, as was also seen in Rodrigues et al.,³⁴ who found an increase in aerobic power in men, from 43.5 to 49.5 ml/kg/min. However, similar to the present study, it was observed a decrease in values from 38.3 to 36.8 ml/kg/min on female, within the age range studied. In turn, 46.0% of students in Canoas (Rio Grande do Sul, Brazil)¹² reached the zone of normality. Both studies corroborate the performance reached by the students from the present sample.

The development of estimative equations of CRF without exercise might be a low-cost alternative to apply in the studies that involve large quantities of subjects, furthermore, it might provide a rapid diagnostic of the current status, mainly in physical education classes or before training program. However, these tools must be used carefully if the purpose is to predict more accurately VO₂max in patients or athletes.³⁵

In the present study it was also possible to identify how BMI exerted influence on VO₂max values in males ($R^2 = 0.850$) and females ($R^2 = 0.907$). This evidence was also observed in Marques-Vidal and Dias³⁶, who studied children in the same age than the present study, Monyeki et al.,³⁷ in 14-year-old adolescents, Castro-Piñeiro et al.,³⁸ in 6- to 17-year-old children and adolescents, Arruda et al.³⁹, in children and adolescents participating in a sports project, and Botelho et al.,⁴⁰ in which the higher the BMI, the lower the cardiorespiratory fitness and physical fitness in general.

It should be emphasized the study of Aires et al.,⁴¹ who evaluated 11- to 18-year-old boys and girls from Portugal. The authors obtained that the students in the range of overweight and obesity had lower levels of CRF, thereby confirming a negative correlation between high fat and VO₂max, furthermore reaffirming that a sedentary lifestyle is a strong predictor of heart disease.

Although cross-sectional studies and the scores of field tests show relative limitation as to its reproducibility, as highlighted in the literature,⁴² they have been widely adopted as benchmarks for health-related physical fitness. Just as the Fitnessgram® tests battery, benchmarks such as Cole et al.⁴³ and World Health Organization (WHO)⁴⁴ play a key role, though Khadilkar et al.¹³ have emphasized the possible constraints in the adoption of their cutoff points due to the anthropometric changes in population over time and the hindering of the categorization of children and adolescents.

The findings of this study were explained to the managers of the Municipal Education from the city of Aracaju, in order to alert them

of the need of the development of public policy towards the adoption of regular physical activities within the school community.

In conclusion, the data of the present study showed that CRF of boys was higher than the girls and there were no differences with respect to BMI. On the other hand, the BF% of the girls showed higher levels. On both sexes there were a minimal prevalence of schoolchildren at risk of overweight and obesity, according to the criteria proposed by Fitnessgram®. Relative to CRF, a high prevalence on both sexes behaved below the area related to health. The linear regression to estimate VO₂max from BMI met expectations for studies on a population scale and confirmed an inverse relationship between both variables, which highlights the need of associate the parameters of physical fitness and health for an early diagnostic of exposure to physical inactivity, obesity and other cardiovascular risk factors.

Conflict of interest

The author declare not have conflict of interest.

References

1. Silva DAS, Petroski EL, Gaya ACA. Anthropometric and physical fitness differences among Brazilian adolescents who practice different team court sports. *J Hum Kinet*. 2013;36:77–86.
2. Erlandson MC, Sherar LB, Mosewich AD, Kowalski KC, Bailey DA, Baxter-Jones ADG. Does the control for biological maturity improve physical activity tracking? *Med Sci Sports Exerc*. 2011;43:800–7.
3. Belli KC, Calegari C, Richter CM, Klafke JZ, Stein R, Viegali PRN. Aptidão cardiorrespiratória de uma amostra regional brasileira distribuída em diferentes tabelas. *Arq Bras Cardiol*. 2012;99:811–7.
4. Maranhão Neto GA, De Leon AP, Lira VA, Farinatti PTV. Assessment of cardiorespiratory fitness without exercise in elderly men with chronic cardiovascular and metabolic diseases. *J Aging Res*. 2012;2012:1–6.
5. Ross WD, Marfell-Jones MJ. Cineantropometria. In: MacDougall JD, Wenger HA, Green HJ, editors. *Evaluación fisiológica del deportista*. 3^{ed}. Barcelona: Editorial Paidotribo; 2005. p. 277–377.
6. Lohman TG. The use of skin fold to estimate body fatness on children and youth. *J Phys Educ Recreat Dance*. 1987;58:98–102.
7. Cyrino ES, Okano AH, Glaner MF, Romanzini M, Gobbo LA, Makoski A, et al. Impact of the use of different skin fold calipers for the analysis of the body composition. *Rev Bras Med Esporte*. 2003;9:150–3.
8. Léger LA, Mercier D, Gadoury C, Lambert J. The multistage 20-m shuttle run test for aerobic fitness. *J Sport Sci*. 1988;6:93–101.
9. Cooper Institute for Aerobics Research. *The prudential FITNESSGRAM test administration manual*. Champaign: Human Kinetics Books; 1999.
10. Liu N, Plowman S, Looney M. The reliability and validity of the 20 m shuttle test in American students 12–15 years old. *Res Q Exerc Sport*. 1992;63:360–5.
11. Duarte MFS, Duarte CR. Validade do teste aeróbio de corrida de vai-e-vem de 20 metros. *Rev Bras Ciên e Mov*. 2001;9:07–14.
12. Bergmann MLA, Halpern R, Bergmann GG. Perfil lipídico, de aptidão cardiorrespiratória e de composição corporal de uma amostra de escolares da 8^a série de Canoas/RS. *Rev Bras Med Esporte*. 2008;14:22–7.
13. Khadilkar VV, Khadilkar AV, Cole TJ, Chiplonkar SA, Pandit D. Overweight and obesity prevalence and body mass index trends in Indian children. *Int J Pediatr Obes*. 2011;6:216–24.
14. Hobold E, Hübner WT. Análise de aspectos morfológicos e funcionais relacionados à saúde dos alunos do projeto “Futuro do judô: iniciação e alto nível através do esporte social”. *Revista Varia Scientia*. 2009;09:11–25.
15. Ronque ERV, Cyrino ES, Mortatti AL, Moreira A, Avelar A, Carvalho FO, et al. Relação entre aptidão cardiorrespiratória e indicadores de adiposidade corporal em adolescentes. *Rev Paul Pediatr*. 2010;28:296–302.
16. Petroski EL, Silva RJS, Pelegrini A. Crescimento físico e estado nutricional de crianças e adolescentes da região de Cotiguiaba, Sergipe. *Rev Paul Pediatr*. 2008;26:206–11.
17. Ara I, Moreno LA, Leiva MT, Gutin B, Casajus JA. Adiposity, physical activity, and physical fitness among children from Aragón, Spain. *Obesity*. 2007;15:1918–24.
18. Malina RM, Bouchard C. *Atividade física do atleta jovem: do crescimento à maturação*. São Paulo: Roca; 2002.
19. Dellagrana RA, Smolarek AC, Laat EF, Campos W. Estado nutricional e desempenho motor de crianças praticantes de handebol. *Fit Perf J*. 2010;9:72–7.
20. Lloyd LK, Bishop PA, Walker JL, Sharp KR, Richardson MT. The influence of body size and composition on FITNESSGRAM® test performance and the adjustment of FITNESSGRAM® test scores for skinfold thickness in youth. *Meas Phys Educ Exerc Sci*. 2003;7:205–26.
21. Short FX, Winnick JP. Test items and standards related to body composition on the Brocicport Physical Fitness Test. *Adapt Phys Act Q*. 2005;22:356–70.

22. Paludo AC, Fernandes RA, Blasquez G, Zambrin LF, Serassuelo Junior H. Concordância entre duas classificações para a aptidão cardiopulmonar em crianças. *Rev Paul Pediatr.* 2012;30:404–8.
23. Bergmann GG, Araújo MLB, Garlipp DC, Lorenzi TDC, Gaya A. Alteração anual no crescimento e na aptidão física relacionada à saúde de escolares. *Rev Bras Cineantropom Desempenho Hum.* 2005;7:55–61.
24. Winsley RJ, Fulford J, Roberts AC, Welsman JR, Armstrong N. Sex difference in peak oxygen uptake in prepubertal children. *J Sci Med Sport.* 2009;12:647–51.
25. Chillón P, Ortega FB, Ruiz JR, De Bourdeaudhuij I, Martínez-Gómez D, Vicente-Rodríguez G, et al. Active commuting and physical activity in adolescents from Europe: results from the HELENA study. *Pediatr Exerc Sci.* 2011;23:207–17.
26. LaMonte MJ, Barlow CE, Jurca R, Kampert JB, Church TS, Blair SN. Cardiorespiratory fitness is inversely associated with the incidence of metabolic syndrome: a prospective study of men and women. *Circulation.* 2005;112:505–12.
27. Ortega FB, Ruiz JR, Castillo MJ, Moreno LA, González-Grossa M, Wärnberg J, et al. Bajo nivel de forma física en los adolescentes españoles. Importancia para la salud cardiovascular futura (Estudio AVENA). *Rev Esp Cardiol.* 2005;58:898–909.
28. Aggio D, Ogunleye AA, Voss C, Sandercock GR. Temporal relationships between screen-time and physical activity with cardiorespiratory fitness in English schoolchildren: a 2-year longitudinal study. *Prev Med.* 2012;55:37–9.
29. Chillón P, Ortega FB, Ferrando JA, Casajus JA. Physical fitness in rural and urban children and adolescents from Spain. *J Sci Med Sport.* 2011;14:417–23.
30. Moliner-Urdiales D, Ruiz JR, Ortega FB, Jiménez-Pavón D, Vicente-Rodríguez G, Rey-López JP, et al. Secular trends in health-related physical fitness in Spanish adolescents: the AVENA and HELENA Studies. *J Sci Med Sport.* 2010;13:584–8.
31. Franchi KMB, Montenegro RM Jr. Atividade física: uma necessidade para a boa saúde na terceira idade. *Rev Bras Promoç Saúde.* 2005;18:152–6.
32. Lee DC, Sui X, Artero EG, Lee IM, Church TS, McAuley PA, et al. Long-term effects of changes in cardiorespiratory fitness and body mass index on all-cause and cardiovascular disease mortality in men. *Circulation.* 2011;124:2483–90.
33. Guedes DP, Miranda Neto JT, Germano JM, Lopes V, Silva AJRM. Health-related physical fitness of schoolchildren: the Fitnessgram program. *Rev Bras Med Esporte.* 2012;18:72–6.
34. Rodrigues AN, Perez AJP, Carletti L, Bissoli NS, Abreu GR. Valores de consumo máximo de oxigênio determinados pelo teste cardiopulmonar em adolescentes: uma proposta de classificação. *J Pediatr.* 2006;82:426–30.
35. Cáceres JMS, Ulbrich AZ, Panigas TF, Benetti M. A non-exercise prediction model for estimation of cardiorespiratory fitness in adults. *Rev Bras Cineantropom Desempenho Hum.* 2012;14:287–95.
36. Marques-Vidal P, Dias CM. Trends in overweight and obesity in Portugal: the National Health Surveys 1995–96 and 1998–99. *Obes Res.* 2005;13:1141–5.
37. Monyeki MA, Neetens R, Moss SJ, Twisk J. The relationship between body composition and physical fitness in 14 year old adolescents residing within the Tlokwe local municipality, South Africa. The PAHL study. *BMC Public Health.* 2012;12:374–81.
38. Castro-Piñero J, Ortega FB, Keating XD, González-Montesinos JL, Sjöström M, Ruiz JR. Percentile values for aerobic performance running/walking field tests in children aged 6–17 years: influence of weight status. *Nutr Hosp.* 2011;26:572–8.
39. Arruda GA, Fernandes RA, Christóforo DGD, Oliveira AR. Relação entre idade cronológica, indicadores de adiposidade corporal e aptidão física relacionada à saúde em meninos e meninas. *Rev Andal Med Deporte.* 2013;6:24–9.
40. Botelho G, Aguiar M, Abrantes C. How critical is the effect of body mass index in physical fitness and physical activity performance in adolescents. *J Phys Educ Sport.* 2013;13:19–26.
41. Aires L, Silva P, Silva G, Santos MP, Ribeiro JC, Mota J. Intensity of physical activity, cardiorespiratory fitness, and body mass index in youth. *J Phys Act Health.* 2010;7:54–9.
42. Guedes DP, Guedes JERP. Manual prático para avaliação em educação física. São Paulo: Manole; 2006.
43. Cole TJ, Bellizzi MC, Flegal KM, Dietz WH. Establishing a standard definition for child overweight and obesity worldwide: international survey. *Br Med J.* 2000;320:1–6.
44. World Health Organization (WHO). Obesity: preventing and managing the global epidemic. Report of a WHO consultation. WHO technical report series 894. Geneva: World Health Organization; 2000.