



Original

ARTÍCULO EN INGLÉS

Continuous training versus interval training in deep water running: health effects for obese women

S.R. Pasetti^a, A. Gonçalves^b and C.R. Padovani^c

^aSchool of Physical Education. University of Campinas (UNICAMP). Campinas, SP, Brasil.

^bSchool of Physical Education. University of Campinas (UNICAMP). Campinas, SP, Brasil.

^cSão Paulo State University. Botucatu. Department of Biostatistics. Campus Rubião Junior, Botucatu, SP, Brasil.

History of the article:

Received August 1, 2011

Accepted December 3, 2011

Keywords:

Aquatic Environment.

Obesity.

Quality of Life.

Palabras clave:

Ambiente acuático.

Obesidad.

Calidad de vida.

ABSTRACT

Objectives. To search intermittent and continuous training (IT and CT, respectively) effects through deep water running for the control and prevention of excessive body fat accumulation and improvement of quality of life.

Methods. Experimental study composed by 30 women, aged between 34 to 58 years old, during 12 weeks, three sessions per week, 47 minutes each. Body composition, cardiorespiratory condition and Quality of Life by WHOQOL-Brief were considered. Student's and Wilcoxon's non parametric tests were applied at 5% significance level.

Results. With the only exception for social domain of quality of life, all investigated variables revealed improvement in both groups for IT in comparison to CT.

Conclusion. Deep water running contributes to body fat reduction, physical fitness evolution and improvement of WHOQOL-Brief domains, regardless of the training type conducted.

© 2012 Revista Andaluza de Medicina del Deporte.

RESUMEN

Entrenamiento continuo vs. interválico corriendo en piscina profunda: efectos sobre la salud de las mujeres obesas

Objetivos. Investigar los efectos del entrenamiento continuo *versus* el entrenamiento intermitente corriendo en piscina profunda para el control y la prevención de la acumulación excesiva de adipocitos sin restricción alimenticia, así como para una mejora del condicionamiento físico y de la calidad de vida.

Métodos. Estudio experimental con treinta mujeres entre 34 y 58 años de edad, con una duración de doce semanas, con tres sesiones semanales de 47 minutos de duración. Se verificó el porcentaje de grasa corporal, la condición cardiorrespiratoria y la calidad de vida mediante la evaluación de criterios del WHOQOL-Abreviado. Se aplicaron la prueba estadística t de Student y la no paramétrica de Wilcoxon. Fue establecido un 5% como nivel de significación.

Resultados. Con la única excepción del dominio social de la calidad de vida, todas las variables investigadas reflejaron una mejora en ambos grupos, con predominio del grupo de entrenamiento intermitente sobre el de entrenamiento continuo.

Conclusión. Independientemente de la forma de entrenamiento utilizada, correr en piscina profunda contribuye a la reducción de la grasa corporal, la evolución de la aptitud física y la mejora de los índices contemplados en el WHOQOL-Abreviado.

© 2012 Revista Andaluza de Medicina del Deporte.

Correspondencia:

S.R. Pasetti.

Rua Renato Reis, 56.

Jardim America, Campinas, SP, Brasil

E-mail: srpasetti@hotmail.com

Introduction

Nowadays, obesity is considered a major public health problem worldwide, as it is increasing at a very fast pace and is associated to diabetes mellitus type II, hypertension, endometrial and colorectal cancer, cardiovascular diseases, sleep apnea, degenerative articular processes and implications on quality of life (QL)¹. Studies show an inverse relationship between the sedentarism rate and QL; the higher the weight, the lower the QL². Systematic physical activities (SPA) cause significant changes in the cardiorespiratory, immunologic, endocrinologic systems, allowing the reduction of body fat, gain of muscle and bone mass, besides preventing chronic degenerative diseases³. Other contributions to QL should also be considered such as reduction of anxiety, improvement of sleep and self-esteem⁴.

Technically, there are two training types for SPA: the continuous (CT) which involves rhythmical exercises with 60-80% of VO_{2max} intensity for extended time⁵ and the one with intervals (IT), which refers to repetition of exercise turns alternated with recovery periods⁶. The first is widely applied as a resource to improve health and to reduce body fat; the second is used for high sports performance. However, IT can be also recommended for those in search of health improvement and reduction of adipocytes concentration. IT results in Vo_{2max} gains and reduction of fat percentage in a more significant way than CT, besides allowing weight maintenance and reduction of cholesterol levels^{6,7}.

The practice of exercises in water environments may be interesting for obese people, as they increase energy consumption when compared to

running training on land, besides reducing injury risks⁸. Among the SPAs performed in water, there is the deep water running (DWR). The practitioner uses a floater attached to the waist allowing the running movements without contacting the pool floor. It is an easy-to-learn activity as the practitioner does not need to know how to swim and there is no impact⁹⁻¹¹.

Considering the SPA benefits for health and QL, the purpose of this study was to investigate the CT and IT effects through SPA for the control and prevention of excessive body fat accumulation and improvement of QL in 30 obese women, aged between 34 and 58 years for 12 weeks, with three 47-minute sessions per week.

Methods

Subjects

Initially, 35 women participated; 30 of them (85.71%) concluded the study in two groups (one with IT comprised 18 volunteers and the other with CT composed of 12) and five were excluded due to excess of absences. Mean age of 46.1±07.87 years (45.33±06.33 years for CT and 46.61±07.88 years for IT). Four (13.3%) were submitted to Menopausal Hormone Therapy, one (3.33%) had diabetes mellitus type II and two (6.66%), mild hypertension. No incidents were registered throughout the intervention.

The selected women met the following criteria: be obese (28 to 35% of body fat), aged between 34-58 years old, having not practiced SPA over the last three months, and have medical authorization or recommendation for physical exercise practice. Training participation




Fig. 1. Periodization scheme of the interval training group.

was not allowed for women who showed: consumptive, hepatic or endocrine disorders, unbalanced renal function or severe dysfunction, chronic obstructive pulmonary disease; pregnancy, neurological or psychiatric problems, abnormal or atypical cardiac response during SPA, use of beta-blockers or anti-obesity drugs, severe hypertension and decompensated diabetes mellitus.

Procedures

The study, characterized as being experimental with two groups, had initial and final assessments and was approved by the Research Ethics Committee of the School of Medicine Science from the University of Campinas (UNICAMP), under the number 640/2005. As dependent variables, physical fitness, reduction of body fat and improvement of QL were considered; the systematic and guided DWR practice was considered as independent variable.

The volunteers' perception of QL was evaluated by using the WHOQOL-Brief (Portuguese version) from the World Health Organization. This tool has 26 standard and closed questions with five response levels, whose answers should be the most appropriate response for the assessments regarding physical, psychological, social relations and environment domains¹². These are considered as a set of subjective perceptions from the participation in the production, circulation and goods consumption processes¹³.

After the WHOQOL-Brief had been filled out, the assessment of the body composition (BC) was started on an individual and systematized basis, which met the following sequence:

- Body mass (kg), the volunteers being weighted in the standing position, barefoot and wearing a bathing suit; and the balance was calibrated before each weighing.
- Skinfold (mm), measured in three consecutive times (tricipital, supra-iliac and mid-thigh) all on the right side¹⁴.

Data obtained in item ii) were used to find body density, a necessary variable to determine body fat percentage (BFP) by using Siri equation¹⁴:

$$\text{Body Density} = 1.099421 - 0.0009929 (X1) + 0.0000023 (X1)^2 - 0.0001392 (X2)$$

$$\%GC = [495 / \text{Body Density}] - 450$$

where X1 is the average sum of the tricipital, suprailiac and mid-thigh skinfolds, and X2, is the individual's age in years.

After these assessments, a period for adaptation to get used to the float and to learn DWR training took place. The training occurs in pool with two meter depth and water in shoulder level. In such conditions the center of gravity is changed when compared to same exercise on land, and is transferred from S1 or S2 sacral vertebra to the thorax (pleural cavity), which renders a more difficult body equilibrium during DWR¹⁵. For this reason, postural and motor re-learning is necessary¹⁶.

This step lasted three sessions, and the last session was dedicated to a specific cardiorespiratory test for DWR¹⁷: whose objective was to estimate the duration time, test heart rate (HR_{peak}) and movements rhythm achieved by the volunteer during the stationary running. It comprises a 4-minute warming with 48 rhythmic elevations of the right leg per minute followed by eleven 2-minute periods without interval. The first starts with 66 elevations per minute which are added by three or four for the others.

The physical conditioning program started after examining the results obtained from the effort test; the prescription for DWR being based on the method proposed by Karvonen¹⁸, which determines the heart rate reserve (HRR).

In each team, the volunteers were allocated to one of the two groups, CT and IT, randomized by simple draw, repeated at each new insertion. This step lasted 12 weeks for both training, totaling 36 sessions.

The sequence for CT group was:

- First week: assessments, adaptation to a liquid environment and DWR learning.
- Second and third week: training performed between 65 and 70% HRR.
- Fourth to sixth week: intensity from 70 to 75% HRR.
- Seventh to ninth week: 75 to 80% HRR.
- Tenth to twelfth week: 80 to 85% of HRR.

For IT, the activities for the first week were the same as for the CT group. Once this phase ended, a sequence with DWR was started at 70-75% HRR associated with the intensive interval training, i.e, high intense stimulus, "sprints", which lasted fifteen seconds with 30 for active recovery, or in terms of weeks (fig. 1):

Table 1

Comparison between mean and standard deviation for continuous and interval training, of fitness variables studied and respective results of statistical test according to assessment times after deep water running practice in obese women

Variable	Continuous Training Group			Interval Training Group		
	Assessment		p value	Assessment		p value
	Initial	Final		Initial	Final	
Body Composition and Anthropometry						
Body mass (kg)	83.86±09.86	82.60±09.67	p<0.05	79.71±12.89	79.05±12.44	p>0.05
Tricipital skinfold (mm)	25.77±04.20	24.44±03.00	p>0.05	24.66±04.53	21.55±04.28	p<0.001
Supra-iliac skinfold (mm)	30.38±04.76	28.05±04.33	p<0.05	29.46±06.56	24.94±00.10	p<0.001
Mid-thigh skinfolds (mm)	33.55±04.32	30.80±03.41	p<0.001	32.00±05.46	28.03±04.06	p<0.001
Sum of skinfolds (mm)	89.69±12.35	83.30±09.57	p<0.001	86.13±14.89	74.59±12.79	p<0.001
Body density (g/cm ³)	1.023±0.007	1.027±0.006	p<0.001	1.025±0.009	1.032±0.008	p<0.001
Body fat percentage	33.96±03.33	32.19±02.65	p<0.001	32.98±03.96	29.63±03.62	p<0.001
Fitness						
Rhythm (# of right leg elevations/min)	68.83±03.18	74.66±07.13	p<0.005	68.55±06.17	75.38±06.72	p<0.001
Test duration (min)	07.83±01.99	11.16±03.76	p<0.005	08.11±02.32	11.77±03.49	p<0.001
HR at rest (bpm)	90.83±08.22	83.75±08.56	p>0.05	88.66±09.88	79.16±10.67	p<0.005
HR at peak (bpm)	143.91±13.60	142.41±17.55	p>0.05	128.50±21.42	128.16±00.65	p<0.05

HR: heart rate.

Table 2
Results of the non-parametric test of domains (%) from the WHOQOL - Brief for the continuous and interval

Descriptive measure	Domains															
	Physical				Psychological				Social Relations				Environment			
	Continuous Training		Interval Training		Continuous Training		Interval Training		Continuous Training		Interval Training		Continuous Training		Interval Training	
I	F	I	F	I	F	I	F	I	F	I	F	I	F	I	F	
Minimum value	39.29	57.14	35.71	60.71	33.33	50.00	25.00	58.33	33.33	50.00	41.67	41.67	34.38	50.00	37.50	46.88
1st Quartile	51.78	62.50	57.14	71.43	52.08	70.83	45.83	70.83	83.33	95.83	58.33	58.33	73.44	79.69	56.25	62.50
Median	57.14	78.57	71.43	80.35	62.50	72.91	62.50	72.91	70.83	75.00	75.00	66.67	60.94	67.19	64.06	70.31
3rd Quartile	71.43	92.86	78.57	89.29	68.75	83.33	70.83	83.33	83.33	95.83	83.33	83.33	73.44	79.69	68.75	78.13
Maximum value	89.29	100.00	89.29	96.43	79.17	91.67	75.00	91.67	91.67	100.0	91.67	100.00	87.50	96.88	81.25	87.50
Average ± SD	61.90±14.87	78.87±16.29	66.07±16.26	80.56±11.76	60.07±13.11	75.00±11.23	58.33±13.85	73.84±9.13	65.97±21.45	79.86±15.67	70.37±14.06	69.90±15.16	63.02±15.53	69.79±14.12	62.15±11.03	68.75±11.69
P value	p<0.005		p<0.001		p<0.001		p<0.001		p<0.005		p<0.05		p<0.05		p<0.05	

I: Initial assessment; F: Final assessment.

- Second to third: eight sprints, allocated in two moments.
- Fourth to sixth: ten sprints, distributed in three sequences.
- Seventh to ninth: twelve at top intensity repetitions, available in three series.
- Tenth to twelfth: fifteen repetitions in three series.

The interval exercises, always considering motion, were distributed into two or three stimuli sets specifically numbered. After checking the initial heart rate (HR_{in}), the evaluated women practiced the exercises in the proposed intensity, i. e. 70-75% HRR for 10 minutes and immediately after they started the first of maximum effort activities, and later returned to the defined target effort.

Two sets of sprints were carried out between the second and sixth training weeks. For other sessions, high intensity stimuli were allocated in three stages, with a five-minute interval between them.

The participants included in the CT and IT groups were monitored during all classes, regarding the heart rate at three-minute intervals, from the end of warm-up up to two minutes after stopping the activity, in to order to follow the effort HR individually established for each session. At the end of the 12-week program, new assessments were made in order to compare the data obtained between the beginning and the end of this study.

Absences were allowed, provided they would not be higher than 20% of total sessions (seven absences); if higher, the volunteer was automatically excluded from the study. For other cases, the classes were replaced following the same schedules of the regular period, and the participant should train observing the session target she missed.

Statistical analysis

As per the analytical assessment, Student's *t* test and Wilcoxon's non-parametric test were applied¹⁹. Data normality was assessed through the Kolmogorov Smirnov test¹⁹. Statistical significance was set at $p < 0.05$ and values were given as mean ± standard deviation²⁰.

Results

Table 1 shows results for BC and physical fitness, considering the initial and final times, according to the training type. The reduction of body mass, in spite of having occurred in CT and IT, was significant for only CT ($p < 0.05$), while the sum of skinfolds ($p < 0.05$) expressed this behavior in both groups.

Body density of $p < 0.001$ indicates improvement in IC and CT; as for the benefits of the continuous and intermittent effort for changes in body composition, it is noticeable that IT has produced twice as much fat reduction as CT ($p < 0.01$).

The increase in rhythm ($p \leq 0.005$) indicates the cardiorespiratory fitness level improved in both groups, being more significant in the IT ($p < 0.001$).

The results in table 2 show the contributions of both training types for improvement of QL DWR contributed similarly for the evolution of women's subjective perception regardless of the proposed interval characteristic.

Discussion

When evaluating the obtained data, one peculiar situation calls the attention related to the compliance with the proposed program, as five

volunteers were excluded due to the excessive number of absences and all belonged to the CT group. According to the participants, the high intensity stimuli were more interesting and motivating than those of the CT, regarded as more monotonous, and this could be the reason for the training interruption.

Considering this, it is well known that exercise practice aiming at weight loss contributes to the favorable evolution of emotional aspects such as humor, self-esteem and body image²¹. In fact, behavior changes towards physical activities and personal relationships were reported in a six-month study on food control and seven weekly 30 to 40-minute long, especially regarding those submitted to high intensity efforts sessions²².

When considering BC, the deep water running practice allowed the reduction of concentrations of localized and general fat, as Assis²³ had already reported in a 15-week study on women with fibromyalgia submitted to stationary DWR.

Body density as well as rhythm were improved, showing respectively increase in body mass²⁴ and cardiorespiratory evolution.

The physical domain in the World Health Organization questionnaire shows improvement and further possibility to be improved with physical activity practice²⁵. The second domain, which involves psychological aspects, includes positive feelings, body image, self-esteem and appearance positively favoring the BC changes²⁶. Probably, people socialization with similar characteristics promoted exchange of experiences and stories with consequent influence on the evolution of social relationships by obese women after DWR²⁷.

Data related to the environment domain call our attention, since they are associated with the essentially economic reality of nature²⁸ as opportunities for recreation and leisure, education, residence, physical environment and satisfaction at work²⁹, on which SPA has little direct interference³⁰. It is believable that such responses are somewhat related with the growing economic trend in Brazil at the time of this intervention. On the other hand, these results were not affected in a similar study conducted by Pasetti; Gonçalves; Padovani¹¹, in a period when the country was beginning to change with a strong social proposal, originating great positive personal expectations.

Studies emphasize that continuous resistance or aerobic exercises are beneficial to QL of different populations as, for instance, those with fibromyalgia²³ or cardiopathy^{31,32} and people with chronic renal problems³³. Data obtained with DWR confirm these findings and indicate that interval training can also contribute to the increase of expressive body fat reduction, besides improving QL of obese women.

In conclusion deep water running has been shown to contribute to body fat reduction, physical fitness evolution and improvement of WHOQOL-Brief domains, regardless of the training type conducted.

References

- Pi-Sunyer FX. Health implications of obesity. *Am J Clin Nutr.* 1991;53:1595S-603S.
- Kolotkin RL, Meter K, Williams GR. Quality of life and obesity. *Obes Rev.* 2001;2:219-29.
- Ghorayeb N, Carvalho T, Lazzoli JK. Non-competitive physical activity for people. In: Ghorayeb N, Barros T, editors. *The exercise: physiological preparation, medical evaluation.* São Paulo: Atheneu; 1999.
- Pasetti SR, Gonçalves A. A for morbid obesity and quality of life: exploring the state of the art. *Health Sci J.* 2006;25:45-51.
- Mcardle WD, Katch FI, Katch VL. *Fundamentals of physiology of exercise.* Rio de Janeiro: Guanabara Koogan; 2002.
- Billat V. *Physiology and methodology of training: from theory to practice.* Barcelona: Editorial Paidotribo; 2002.
- Yoshioka M, Doucet E, St-Pierre S, Alméras N, Richard D, Labrie A, et al. Impact of high-intensity exercise on energy expenditure, lipid oxidation and body fatness. *Int J Obes Relat Metab Disord.* 2001;25:332-9.
- Gutin B, Barbeau P, Owens S, Lemmon CR, Bauman M, Allison J, et al. Effects of exercise intensity on cardiovascular fitness, total body composition, and visceral adiposity of obese adolescents. *Am J Clin Nutr.* 2002;75:818-26.
- Fernández AC, Mello MT, Tufik S, Castro PM, Fisberg M. Influence of aerobic and anaerobic training on body fat mass in obese adolescents. *Braz Sports Med J.* 2004;10:152-8.
- Cureton KJ. Physiological responses to water exercise. In: Ruoti RG, Morris DM, Cole A, editors. *Aquatic rehabilitation.* Philadelphia, PA: Lippincott-Raven; 1997.
- Pasetti SR, Gonçalves A, Padovani CR. Deep water running to improve the physical fitness of middle-aged obese women: an experimental study of a single group. *Braz J Phys Educ Sports.* 2006;20:297-304.
- Development of the World Health Organization WHOQOL-Brief quality of life assessment. The WHOQOL Group. *Psychol Med.* 1998;28:551-8.
- Gonçalves A. A Presentation. In: Gonçalves A, Vilarta R, editors. *Quality of life and physical activity: exploring theory and practice.* São Paulo: Manole; 2004.
- Jackson AS, Pollock ML, Ward A. Generalized equations for predicting body density of women. *Med Sci Sports Exerc.* 1980;12:175-81.
- Bates A, Hanson N. *Aquatic exercise therapy.* Philadelphia: WB Saunders; 1996.
- Galdi EH, Goncalves A, Vilarta R, Clemente JF, Vicentin AP, Leite JP. *Learning to swim with the university.* Campinas, São Paulo: IPES; 2004.
- Wilder RP, Brennan DK, Schotte DE. A standard measure for exercise prescription for aqua running. *Am J Sports Med.* 1993;21:45-8.
- Karvonen MJ, Kentala E, Mustala O. The effects of training heart rate: a longitudinal study. *Ann Med Exp Biol Fenn.* 1957;35:307-15.
- Norman GR, Streiner DL. *Biostatistics: the bare essentials.* Saint Louis: Mosby Book; 1994.
- Gonçalves A. The testing of hypotheses and validation of instrumental interpretation (statistical inference) In: Marcondes M, Lakatos EM, editors. *Research Techniques.* São Paulo: Atlas; 1982.
- Blissmer B, Riebe D, Dye G, Ruggiero L, Greene G, Caldwell M. Health-related quality of life following a clinical weight loss intervention among overweight and obese adults: intervention and 24 month follow-up effects. *Health Qual Life Outcomes.* 2006;4:43.
- Gallagher KI, Jakicic JM, Napolitano MA, Marcus BH. Psychosocial factors related to physical activity and weight loss in overweight women. *Med Sci Sports Exerc.* 2006;38:971-80.
- Assis MR, Silva LE, Alves AM, Pessanha AP, Valim V, Feldman D, et al. A randomized controlled trial of deep water running: clinical effectiveness of aquatic exercise to treat fibromyalgia. *Arthritis Rheum.* 2006;55:57-65.
- Perna F, Bryner R, Donley D, Kolar M, Hornsby G, Sauer J, et al. Effect of diet and exercise on quality of life and fitness parameters among obese individuals. *Journal of exercise physiology.* 1999;2. [Access 2009 Jun 15]. Available from: <http://faculty.css.edu/tboone2/asep/april199B.htm>
- Fontaine KR, Barofsky I, Bartlett SJ, Franckowiak SC, Andersen RE. Weight loss and health-related quality of life: results at 1-year follow-up. *Eat Behav.* 2004;5:85-8.
- Becker BE. Physical principles of water In: Ruoti RG, Morris DM, Cole A, editors. *Aquatic Rehabilitation.* São Paulo: Manole; 2000. p.17-28.
- Gonçalves A, Moura AC, Vicentin AP, Leo CC, Del Vecchio FB, Mantellini GG, et al. Public health and physical activity: clippings of a current performance. *Connections.* 2006;4:1-14.
- Fleck MP, Louzada S, Xavier M, Chachamovich E, Vieira G, Santos L, et al. Application of the Portuguese version of the abbreviated instrument of quality of life "WHOQOL-bref". *Rev Saude Publica.* 2000;34:178-83.
- WHO. World Health Organization, Mental Health Division. WHOQOL-Abreviado. Portuguese version of tools for quality of life evaluation of World Health Organization (WHOQOL) 1998, developed by WHOQOL Center for Brazil. Department of Psychiatry and Legal Medicine of the Federal University of Rio Grande do Sul. [Access 2005 Jun 16]. Available from: <http://www.ufrgs.br/psiq/whoqol84.html>
- Glise H, Wiklund I. Health-related quality of life and gastrointestinal disease. *J Gastroenterol Hepatol.* 2002;17 Suppl:S72-84.
- Bocalini DS, Dos Santos L, Serra AJ. Physical exercise improves the functional capacity and quality of life in patients with heart failure. *Clinics (Sao Paulo).* 2008;63:437-42.
- Winter MM, Bouma BJ, Van Dijk AP, Groenink M, Nieuwkerk PT, Van der Plas MN, et al. Relation of physical activity, cardiac function, exercise capacity, and quality of life in patients with a systemic right ventricle. *Am J Cardiol.* 2008;102:1258-62.
- Reboredo MM, Henrique DM, Bastos MG, De Paula RB. Physical exercise in dialysis patients. *Braz Spor Med Journal.* 2007;13:427-30.