



Original

ARTÍCULO EN INGLÉS

Nutritional analysis of diet at base camp of a seven thousand-metre mountain in the Himalayas

M. Mariscal-Arcas^a, C. Carvajal^a, C. Monteagudo^a, J. Lahtinen^d, M. C. Fernández de Alba^b, B. Feriche^c and F. Olea-Serrano^a

^aNutrition and Food Science Department. University of Granada. Spain.

^bAndalusian Centre of Sport Medicine (CAMD). Junta de Andalucía. Spain.

^cPhysical Education and Sport Department. University of Granada. Spain.

^dAltitude Training Centre (CAR) of Sierra-Neveda. Consejo Superior de Deportes (CSD). Spain.

History of the article:

Received April 5, 2010

Accepted July 2, 2010

Key words:

Mountaineer.

Diet.

Carbohydrate.

Fats.

Palabras clave:

Alpinista.

Dieta.

Carbohidratos.

Grasas.

ABSTRACT

Objective. To evaluate the diet of a group of high-mountain climbers at 4,500 metres.

Methods. A descriptive cross-sectional study was performed on the diet of a group of high-mountain climbers at their base camp (4,500 m).

Results. The mean intake was 11.85 MJ/day (2,833 kcal/day), which provided an inadequate supply of energy and micronutrients to replenish deposits. Their mean carbohydrate intake (39.5%) was excessively low, since carbohydrate-rich diet favours acclimatization and the capacity for recovery. Their daily intake of 1.5-2.5 g/protein/kg of bodyweight was very similar to recommendations (1.5-2.0 g/kg/day).

Conclusions. The climbers underwent a drastic change from their habitual Mediterranean diet, rich in monounsaturated fats (largely olive oil), to a diet rich in polyunsaturated fats (largely soy oil). The Sherpa-prepared diet on this expedition was not balanced. It was rich in saturated polyunsaturated fats and relatively poor in proteins and especially carbohydrates, similar to the traditional diet of climbers. Dietary strategies need to be developed to increase the intake of nutrients that favours the physical recovery of climbers and their altitude-acclimatization and to avoid micronutrient deficiencies.

© 2010 Revista Andaluza de Medicina del Deporte.

RESUMEN

Análisis nutricional de la dieta en un campamento base a 7.000 metros en una montaña del Himalaya

Objetivo. Evaluación de la dieta de un grupo de escaladores de alta montaña a 4.500 metros.

Método. Estudio descriptivo de corte transversal realizado sobre la dieta de un grupo de escaladores de alta montaña en su campamento base (4.500 m).

Resultados. La ingesta media fue de 11,85 MJ/día (2.833 kcal/día), que proporcionan un suministro inadecuado de energía y micronutrientes para reponer los depósitos. El consumo promedio de los hidratos de carbono (39,5%) era excesivamente bajo, ya que la aclimatación y la capacidad de recuperación se favorece con una dieta rica en carbohidratos. La ingesta diaria de 1,5-2,5 g/proteína/kg de peso corporal fue muy similar a las recomendaciones (1,5-2,0 g/kg/día).

Conclusiones. Los escaladores sufrieron un cambio drástico de su dieta mediterránea habitual, rica en grasas monoinsaturadas (aceite de oliva en gran medida), al pasar a una dieta rica en grasas poliinsaturadas (principalmente aceite de soja). La dieta preparada por los serpas, en esta expedición, no era equilibrada. Era una dieta rica sobre todo en grasas saturadas y poliinsaturadas, relativamente pobre en proteínas y carbohidratos, similar a la dieta tradicional de los escaladores. Se deben desarrollar estrategias en la dieta para aumentar la ingesta de nutrientes que favorezcan la recuperación física de los escaladores y su aclimatación a la altura y así evitar carencias de micronutrientes.

© 2010 Revista Andaluza de Medicina del Deporte.

Correspondence:

F. Olea-Serrano.

Nutrition and Food Science Department.

University of Granada.

Campus de Cartuja s/n

18071 Granada. Spain.

E-mail: folea@ugr.es

Introduction

Nutrition plays an important role in the achievement of optimal performance in sports and is especially critical in resistance sports. The resistance sport of alpinism combines high physical exercise with hypoxia-induced anorexia, especially at altitudes above 6,000 metres. Studies have shown that the organism can oxidize fats and carbohydrates in a normal manner up to 5,000 metres, therefore any weight loss below this altitude can be attributed to inadequate calorie intake¹⁻³. Weight loss above 5,000 metres appears inevitable due to the loss of appetite and nausea produced by altitude sickness and the metabolic changes required to obtain energy.

It has generally been accepted that the human diet should contain 55-65% carbohydrates, 12-20% proteins and 20-30% fats⁴, although the WHO recently increased the recommended proportion of carbohydrates to 75%⁵. However, some authors have supported increasing the protein intake at the expense of fats at high altitudes, with each component comprising 15-20% of the diet. They propose a daily intake of at least 7 g of carbohydrates per kg of weight (3.2 g per pound) plus 1.5-2.0 g of protein, with remaining calories consumed as fats⁶. Special caution should be taken with traditional fat-rich diets, which can lead to chronic muscle fatigue; they provide an inadequate amount of readily available carbohydrates, and diets rich in fatty acids require more oxygen during metabolism, slowing acclimatization. In a hypobaric chamber study⁷, in which altitude hypoxia was simulated by the gradual decompression of the chamber, subjects consumed *ad libitum* from a range of appetizing dishes but lost weight in proportion to the simulated altitude *versus* controls offered the same food under normal pressure conditions. Therefore, hypobaric hypoxia appears to be sufficient to produce weight loss, probably due to anorexia and poor intestinal absorption. These effects are increased in real expeditions^{8,9}.

Oxygen saturation values of around 70% are essential for good absorption in the small intestine and avoidance of weight loss¹⁰⁻¹¹. Roberts et al¹² suggested that acclimatized individuals may have lower fat metabolism activity at high altitudes, influencing carbohydrate availability. On the other hand, McClelland et al¹³ concluded that the relative carbohydrate contribution does not change after altitude acclimatization and that the metabolic utilization of fuel is mainly influenced by the relative intensity of exercise, as at sea level.

There is little evidence that chronic or acute exposure to altitude increases the demand for any specific micronutrient, although there have been reports that iron and vitamin E supplements may be beneficial¹⁴. Studies of the effects of cold, energy expenditure, ultraviolet ray exposure and atmospheric composition indicate that supplementation with certain vitamins has a highly desirable antioxidant function in the high mountains¹⁵⁻¹⁸. At high altitudes, the prolonged consumption of vitamins with antioxidant properties can prevent the slowing of blood flow and the reduction in physical function associated with free-radical induced damage^{19,20}. Dietary treatments that preserve or enhance the fluidity or deformability of the red blood cell membrane improve the transfer of oxygen to tissues, and action to improve oxygen delivery to tissues under hypoxia conditions is generally advantageous for the functioning of the organism. Exposure to hypoxia and peroxidation of unsaturated fatty acids of the red blood cell membrane reduce its deformability and its capacity to adapt to capillaries. Improved membrane fluidity (increased deformation capacity) can be achieved by two dietary mechanisms, for instance, diet supplementation with polyunsaturated fatty acids

or antioxidants (for instance, vitamin E), in order to protect polyunsaturated fatty acids in membranes from peroxidation by free radicals.

Iron supplements can also be beneficial in this situation. According to observations of the response of erythropoiesis and haemoglobin synthesis at high altitudes, a normal iron intake appears adequate for males, whereas women may benefit from iron supplementation. The intake of oral iron supplements (ferrous sulphate, 200-300 mg/day) has been proposed during 2-3 weeks before a climb and during 2-4 weeks while at altitude, with the caveat that an increase in free radicals may be produced²¹⁻²³.

In November and December 2002, seven climbers from Andalusia (Southern Spain) attempted the ascent of Jannu ("sleeping giant" in Nepalese), a Himalayan mountain of 7,710 meters in the Kangchenjunga region of Nepal.

The objective of the present study was to evaluate the diet of a group of high-mountain climbers at 4,500 metres, analyzing the types and amounts of nutrients.

Methods

Subjects

A study was undertaken of the diet followed by seven male climbers on the "Andalusia K2 2003" expedition; they were aged from 27 to 42 years and all had considerable mountaineering experience. Before acceptance as expedition member, each was informed both verbally and in writing of the full extent of the proposed research, and written informed consent was then obtained from each subject. All were in excellent physical condition prior to departure. Table 1 lists the responsibilities of expedition members and their anthropometrical parameters, calculated according to FAO/WHO²⁴.

Procedure

The expedition comprised two phases: firstly ascent of Jannu in October, November, and December 2002; and secondly ascent of K2 (at 8,611 meters, the second highest mountain in the world) in June, July, and August 2003. The expedition was organized by the Andalusian Mountaineering Federation and subsidized by the Department of Tourism and Sports of the Andalusian Regional Government. The investigation was approved by the ethics committee of the University of Granada and all subject data were coded to maintain confidentiality.

Food and supplements

A register was pre-prepared for recording all data related to nutrition throughout the expedition. We recorded and analysed all of the food consumed at base camp during a 17-day period. Food items were brought from Katmandu and prepared by two Sherpa cooks.

Four daily meals were prepared: breakfast, mid-morning snack, lunch, and evening meal. Varied menus were provided during the 17-day study period, prepared from both fresh and preserved foods. The amounts were calculated from the weight of the food items, measured by the climber responsible for food. They were weighed by means of a kitchen scale of 2 kg capacity with 20 g increments (Hanson UK Ltd.

Table 1
Responsibilities and characteristics of expedition members

| | Leader | Sub-Leader | Technical material | Physician | Audiovisual materials | Communications | Food | Mean |
|----------------------------------|--------|------------|--------------------|-----------|-----------------------|----------------|-------|-------|
| Age (years) | 37.00 | 42.00 | 30.00 | 38.00 | 36.00 | 39.00 | 27.00 | 35.57 |
| Initial weight (kg) | 65.20 | 68.30 | 71.30 | 72.00 | 66.00 | 80.40 | 71.10 | 70.61 |
| Final weight (kg) | 62.00 | 65.10 | 67.30 | 70.30 | 63.20 | 75.40 | 66.20 | 67.07 |
| Weight difference (kg) | 3.20 | 3.20 | 4.00 | 1.70 | 2.80 | 5.00 | 4.90 | 3.54 |
| Height (m) | 1.71 | 1.72 | 1.78 | 1.80 | 1.72 | 1.80 | 1.74 | 1.75 |
| Initial BMI (kg/m ²) | 22.23 | 22.98 | 22.41 | 22.22 | 22.31 | 20.83 | 23.45 | 22.35 |
| BMR (MJ)* | 6.77 | 6.92 | 7.10 | 7.11 | 6.82 | 7.49 | 7.37 | 7.08 |

*Calculated according to FAO/WHO, 2001 (18-30 years old, 0.063 kg + 2.896; 30-60 years old, 0.048 kg + 3.653). Mean weight loss = 3.57 kg. BMR: basal metabolic rate.

Table 2
Description of the foods consumed at base

| Vegetables | Consumption (g and g/day) | Frequency during 17 days | Protein foods | Consumption (g and g/day) | Frequency during 17 days | Fruit | Consumption (g and g/day) | Frequency during 17 days |
|----------------------|---------------------------|--------------------------|---------------------------|---------------------------|--------------------------|--------------------------------|---------------------------|--------------------------|
| Carrot purée | 300 (75) | 4 | Egg | 1,200 (150) | 8 | Preserved compote | 1,440 (180) | 8 |
| Carrot | 847 (60) | 14 | Lean yak steak | 1,500 (100) | 15 | Peach in syrup | 1,080 (180) | 6 |
| Cabbage | 375 (75) | 5 | Canned sardines | 125 | 1 | Preserved mango | 600 (200) | 3 |
| Boiled potatoes | 800 (115) | 7 | Frankfurt sausage | 325 (65) | 5 | Sugar | 140 (10) | 14 |
| Preserved sweet corn | 50 | 1 | Tuna in pastry | 200 | 1 | Honey | 20 | 1 |
| Endives | 180 (90) | 2 | Breaded chicken | 120 | 1 | Homemade chocolate cake | 150 | 1 |
| Cauliflower | 75 | 1 | Potato omelet | 1,350 (150) | 9 | Cinnamon sponge cake | 150 | 1 |
| Cauliflower cheese | 300 (75) | 4 | Powdered whole cow's milk | 1,800 (165) | 11 | Custard | 250 (125) | 2 |
| Mushrooms | 470 (94) | 5 | Yak cheese | 180 (20) | 9 | Jam | 25 | 1 |
| Peas | 240 (80) | 3 | Carbohydrate foods | | | Wheat bread | 100 | 1 |
| Onion, garlic | 150 (30) | 5 | White rice | 350 (87.5) | 4 | Other food | | |
| Vegetable soup | 250 | 1 | Corn flour | 165 (55) | 3 | Mayonnaise | 40 | 1 |
| Green beans | 120 (40) | 3 | Cornflakes | 180 (30) | 6 | Packet soups | 200 (25) | 8 |
| Lettuce | 200 (50) | 4 | Chapatti with egg | 575 (115) | 5 | Pizza | 550 (275) | 2 |
| Corn | 1,400 (100) | 14 | Popcorn | 500 (100) | 5 | Tea | 4,250 (250) | 17 |
| Chickpeas | 160 (80) | 2 | Pasta | 480 (80) | 6 | Spices (pepper, mustard, etc.) | Ad libitum | |
| Tomato sauce | 660 (55) | 12 | Pasta with vegetables | 640 (80) | 8 | | | |
| | | | Chapattis | 100 | 1 | Fats | | |
| | | | Potato chips | 100 | 1 | Butter | 20 | 1 |
| | | | Spaghetti in white sauce | 80 | 1 | Soy oil | Ad libitum | 17 |

ref. 205 H, 700320363 Rev. 6.11). The same climber maintained the register, in which all foods purchased were noted and the food consumed on each day was recorded. The diet was analyzed by using the Dietsource software package²⁵. When necessary, information on the composition of local foods (for instance, yak meat) was added to the database of this program, derived from the Food Composition Table for use in East Asia²⁶. The energy intake (EI) was estimated from the daily menus. All mountaineers ate exactly the same meals and always ate all of the food offered, therefore the daily EI can be assumed to be identical for all study subjects. The estimated-EI value was related to the total energy expenditure (TEE) according to equations proposed by FAO/WHO²⁴ ($TEE = 662 - 9.53 \times \text{age (years)} + PA \times 15.91 \times \text{weight (kg)} + 539.6 \times \text{height (m)}$), taking account of the basal metabolic rate (BMR) and physical activity level (PAL) of the group²⁷. TEE values were calculated considering active PAL values (1.70-1.99) or, on climbing days, vigorous PAL values (2.00-2.40) PAL values.

Statistical analysis

Mean, minimum and maximum levels of nutrients in the menus were calculated, and the Student's t-test was used to compare nutritional data gathered during the 17 days of the expedition (estimated-EI) with nutrient intake recommendations (2005 Dietary Reference Intakes [DRI] data) for this population²⁸. The normal distribution of nutrient values was checked and confirmed by using Kolmogorov-Smirnov and Shapiro-Wilke tests^{29,30}. $P < 0.05$ was considered statistically significant. SPSS version 15.0 software was used for all data analyses.

Results

Table 2 lists food groups in the diet, amounts per person and frequency of their consumption. Tables 3 and 4 show results of the nutritional analysis of the meals consumed at the base camp during the 17-day

Table 3

Nutrients and energy provided by the diet and mean distribution of fats in studied diet

| Nutrients/day | Minimum | Maximum | Mean | SD | % energy |
|-----------------------|---------|----------|----------|--------|----------|
| Proteins (g) | 22.30 | 167.10 | 110.70 | 46.20 | 15.50 |
| Fats (g) | 60.90 | 206.10 | 142.10 | 47.60 | 45.50 |
| Carbohydrates (g) | 94.10 | 399.90 | 277.60 | 84.80 | 39.50 |
| Fiber (g) | 3.90 | 20.00 | 12.60 | 4.80 | |
| Energy (MJ) | 4.83 | 16.70 | 11.85 | 3.77 | 100.00 |
| Saturated fats (g) | 8.90 | 73.10 | 38.00 | 20.30 | 33.60* |
| Monosaturated fat (g) | 19.70 | 37.40 | 25.50 | 7.50 | 27.60* |
| Polysaturated fat (g) | 20.20 | 77.70 | 49.40 | 5.20 | 43.70* |
| Cholesterol (mg) | 104.00 | 2,164.00 | 1,051.70 | 618.90 | |

*% energy from 45.5 fats energy.

SD: standard deviation.

study and the comparison (Student's t-test for one sample) with DRI values. Minimal estimated-EI (calculated from mean portions consumed by the group) and hence minimal estimated-EI/BMR ratios were observed on days 1 and 11 of the base camp stay. The mean estimated-EI/BMR (SD) of the group was 1.67 (0.53), ranging from 0.64 to 2.47. Table 5 shows the comparison between the TEE values calculated for each climber (according to weight, age and physical activity) and their estimated-EI values (see table 5 footnotes). When active PAL values were considered, no differences between mean estimated-EI and TEE values were found, but when vigorous PAL values were assumed, significant differences ($P < 0.05$) were found for five of the climbers.

Climbers were weighed at the start and end of the entire 2-month expedition, finding weight losses ranging from 2 to 5 kg (table 1).

Discussion

On a high-mountain climb, 15.90–25.10 MJ (3,800–6,000 kcal) are expended daily according to the type of physical activity involved¹¹. It is evidently important to consume the amount of calories required by the organism. Climbers lose at least 3% of their body weight after eight days at 4,300 meters and 15% after three months at 5,300–8,000 meters³¹. This loss can produce a reduction in mental and physical functioning. Some studies have suggested that the intake of carbohydrates at altitude

improves blood oxygenation and provides a more efficient energy source compared with fats or protein². The macronutrient and energy levels observed in the present climbers differed widely from the usual recommendations for climbs above 6,000 meters. Nevertheless, no significant differences were found between the mean estimated-EI values estimated from the diet (11.85 MJ/day [2,832.5 kcal/day]) and the TEE values calculated for each climber according to 2001 FAO/WHO recommendations²⁴ for adults with moderate activity. When the TEE was calculated for adults engaged in vigorous activity (PAL = 2.00–2.40), however, a significant difference with the estimated-EI value was found for five climbers. It should be borne in mind that this level of activity cannot be maintained during long time periods.

Among the 17 acclimatization days studied, estimated-EI/BMR values of > 2.00 were recorded on the only 7 days when they were climbing, an intake of at least 8.37 MJ/day (2,000 kcal/day) is recommended during long-duration activities to avoid iron or calcium deficits, depletion of glycogen deposits and dehydration³¹. Since these climbers were at 4,500 m and the aim of their base camp stay was to replenish energy deposits, their mean intake of 11.85 MJ/day (2,833 kcal/day) provided an inadequate supply of energy and micronutrients. These observations are similar to those reported by most studies, either in hypobaric chamber or *in situ*³.

The production of energy per litre of oxygen is higher when carbohydrates are the energy source, regardless of the oxygen pressure of inhaled air. Carbohydrates are also a more efficient energy source for working under conditions of reduced oxygen pressure³². The 39.5% carbohydrate intake of the present study group was very low, since around 55–65% is usually recommended to avoid muscle exhaustion from excessive depletion of glycogen reserves and to maintain adequate glycaemia against excessive protein catabolism. In short, a carbohydrate-rich diet favours acclimatization and the capacity for recovery^{4,31}.

The climbers had a daily intake of 1.5–2.5 g of proteins per kg of bodyweight, very similar to the recommendations of most authors consulted (1.5–2.0 g/kg/day)^{33,34} and representing 12–15% of the total energy consumed³⁵. Nevertheless, some authors have proposed a higher intake of 2.5–3 g/kg/day³⁶. An adequate intake of proteins and glucose liquids is essential to prevent excessive weight loss under conditions of very high catabolism. Furthermore, prolonged exercise has a similar immunosuppressive effect to that of diets deficient in proteins and specific micronutrients³⁷. Although no amino acid supplements were

Table 4

Mean (total/daily) intake of micronutrients during the 17 days at base camp

| Nutrients intake/day | Minimum | Maximum | Mean (SD) | DRI | T Test | P |
|-------------------------|---------|---------|-------------------|-------|---------|--------|
| Phosphorus (mg/day) | 474.6 | 3,027.1 | 1,949.2 (884.8) | 700 | 5.647 | 0.0001 |
| Magnesium (mg/day) | 163.5 | 485.3 | 342.0 (108.2) | 420 | -2.881 | 0.011 |
| Calcium (mg/day) | 142.1 | 2,774.5 | 1,487.4 (1,019.0) | 1,000 | 1.913 | 0.075 |
| Iron (mg/day) | 5.2 | 29.1 | 17.1 (6.3) | 8 | 5.767 | 0.0001 |
| Zinc (mg/day) | 4.5 | 25.1 | 15.5 (7.4) | 11 | 2.441 | 0.028 |
| Iodine (ug/day) | 7.0 | 101.7 | 47.3 (25.1) | 150 | -16.303 | 0.000 |
| Selenium (ug/day) | 6.5 | 78.8 | 38.5 (23.4) | 55 | -2.811 | 0.013 |
| Ascorbic acid (mg/day) | 18.3 | 277.1 | 89.5 (73.8) | 90 | -0.025 | 0.980 |
| Thiamin (mg/day) | 0.5 | 2.4 | 1.5 (0.6) | 1.2 | 2.540 | 0.023 |
| Riboflavin (mg/day) | 0.5 | 6.8 | 3.0 (1.9) | 1.3 | 3.648 | 0.002 |
| Nicotinic acid (mg/day) | 3.6 | 32.1 | 19.9 (8.7) | 16 | 1.815 | 0.090 |
| Pyridoxine (mg/day) | 0.6 | 4.7 | 2.7 (1.6) | 1.3 | 3.678 | 0.002 |
| Folic acid (ug/day) | 77.4 | 302.3 | 168.4 (66.8) | 400 | -13.852 | 0.0001 |
| Cyanocobalamin (ug/day) | 1.1 | 39.9 | 8.1 (8.9) | 02.4 | 2.542 | 0.023 |
| Vitamine E (mg/day) | 12.8 | 24.5 | 18.8 (4.3) | 15 | 3.55 | 0.003 |

DRI: Dietary Reference Intakes, 2005.

SD: standard deviation.

Table 5
Comparison of means of EI (MJ/day) vs. TEE (MJ/day)

| Climber number | TEE (active) (MJ/d) | T Test | P | TEE (vigorous) (MJ/d) | T Test | P |
|----------------|---------------------|--------|-------|-----------------------|--------|-------|
| 1 | 11.53 | 0.342 | 0.737 | 13.41 | -1.653 | 0.119 |
| 2 | 11.61 | 0.259 | 0.799 | 13.54 | -1.791 | 0.094 |
| 3 | 12.50 | -0.692 | 0.499 | 14.52 | -2.823 | 0.013 |
| 4 | 12.33 | -0.502 | 0.623 | 14.36 | -2.661 | 0.018 |
| 5 | 11.68 | 0.181 | 0.858 | 13.58 | -1.836 | 0.086 |
| 6 | 12.95 | -1.165 | 0.262 | 15.11 | -3.453 | 0.004 |
| 7 | 12.51 | -0.699 | 0.495 | 14.50 | -2.808 | 0.013 |

Equation used in the calculation: *TEE = 662-9.53 × age (years) + PA × 15.91 × weight (kg) + 539.6 × height (m). PA = 1.25 if PAL is considered to be 1.70-1.99 (active); PA = 1.48 if PAL is considered to be 2-2.4 (vigorous).

PA: physical activity; TEE: total energy expenditure.

used in this expedition, some authors have proposed that valine, leucine and isoleucine supplementation prevents loss of muscle mass during acute hypobaric hypoxia³⁸. In fact, the intake of these amino acids by the present climbers was similar to levels recommended for the general population. Supplementation with probiotics and glutamine appears to play a major role in immunofunction³⁹. With regard to the origin of the protein, some studies have indicated that vegetable proteins produce a small reduction in free fatty acids⁴⁰, which could be beneficial in diets that are very rich in fats.

In contrast to their carbohydrate content, the fat levels of the present diet were excessively elevated (45%). According to the American Dietetic Association and the American College of Sports Medicine³¹, fats should constitute at least 15% but not more than 20-25% of a diet. As mentioned above, fats are not the most efficient energy source under hypoxia conditions. The overconsumption of fats is a frequent observation, both among the general population and in different sports settings, for instance, swimming⁴¹.

Diets that are very rich in polyunsaturated fatty acids, especially linoleic acid, appear to reduce the absorption and utilization of iron, zinc and magnesium, reducing sports performance⁴². Nevertheless, some authors concluded that the role of linoleic acid remains unclear³⁹. It should be taken into account that the present climbers underwent a drastic change in their diet. They all live in a Mediterranean region where fat consumption is largely in the form of olive oil, rich in monounsaturated fats (oleic acid), whereas the fats they consumed during the expedition were usually in the form of soy oil, rich in linoleic acid. Soy oil contains 45-60% linoleic acid, whereas olive oil contains 61-82% oleic acid, which presents greater resistance to auto-oxidation processes and the formation of free radicals^{43,44}. If the climbers had used olive oil instead of soy oil, their diet would have contained a much higher proportion of monounsaturated versus polyunsaturated fats.

The well-documented importance of iron, calcium, magnesium and vitamins D and C is even greater in situations of high oxidative stress, such as those produced in high mountains and by adaptation to high altitudes^{45,46}. The increased excretion of iron, copper and manganese in intense and prolonged exercise was reported to produce a negative mineral balance^{42,47}, although Fogelholm³⁹ claimed neither the mineral nor vitamin balance is usually a problem for athletes, with the exception of iron and calcium in females. The supply of iron appears to have been adequate in the present study group (mean intake of 17.1 mg/day). Several authors have recommended that iron supplements may be taken up to two to three weeks before an expedition, maximizing iron levels, but once in the mountain they should only be supplied by the daily food intake,

thereby avoiding the release of free radicals produced by iron supplements in hypoxia situations^{48,49}. Excessive micronutrient supplementation (mainly vitamin C, B-group vitamins, and iron) during climbs has been contraindicated, since it does not improve sports performance and may have undesirable effects⁵⁰. Nevertheless, vitamin C and general vitamin supplementation appears to be appropriate after exercise^{51,52}.

The diet under study showed some interesting qualitative differences with the usual traditional Mediterranean diet⁵³ of the climbers. It included novel foods such as yak products (meat, butter and cheese) and soy oil instead of olive oil, as mentioned above, and there was an absence or relative scarcity of fish, fruit, and vegetables (table 2).

The nutrient composition of the diet of these mountain climbers was not ideal. It was rich in saturated fats of animal origin and polyunsaturated fats from soy oil and was relatively poor in proteins and especially in carbohydrates, similar to the traditional diets of climbers. This type of nutrition does not favour acclimatization, because metabolism of the excessive fats requires greater supplies of oxygen at the expense of other tissues. An inadequate carbohydrate intake compromises effective muscle and liver recovery, produces a hyperglycaemia that increases protein catabolism, and accelerates weight loss⁵⁴. The diet had a very low content of antioxidant vitamins, reducing the climbers' defences against free radicals and increasing the risk of damage to cell membranes. In conclusion, the diet of these climbers was not appropriate under these conditions. The lessons drawn from this study will serve to improve the nutritional intake of this team of alpinists during future expeditions.

Conflict of interest statement

The authors declare that there are no conflicts of interest.

Acknowledgements

The authors wish to thank Richard Davies for his assistance with the English version. This study was supported by the Junta de Andalucía, Spain (Research Group AGR-255 "Nutrition, Diet and Risk Assessment") and a postdoctoral grant from the University of Granada, Spain.

References

- Zamboni M, Armellini F, Turcato E, Robbi R, Micciolo R, Todesco T, et al. Effect of altitude on body composition during mountaineering expeditions: interrelationships with changes in dietary habits. *Ann Nutr Metab*. 1996;40(6):315-24.
- Brouns F. Nutritional aspects of health and performance at lowland and altitude. *Int J Sports Med*. 1992;13 Suppl 1:S100-6.
- Westerterp KR. Energy and water balance at high altitude. *News Physiol Sci*. 2001;16:134-7.
- Butterfield GE. Nutrient requirements at high altitude. *Clin Sports Med*. 1999;18:607-21.
- FAO/WHO Diet, Nutrition and the Prevention of Chronic Diseases. Joint FAO/WHO Expert Consultation Geneva, 2003.
- Rodríguez NR, Di Marco NM, Langley S. American Dietetic Association; Dietitians of Canada; American College of Sports Medicine, American College of Sports Medicine position stand. Nutrition and athletic performance. *Med Sci Sports Exerc*. 2009;41(3):709-31.
- Rose MS, Houston CS, Fulco CS, Coates G, Sutton JR, Cymerman A, et al. Operation Everest. II: Nutrition and body composition. *J Appl Physiol*. 1988;65:2545-51.
- Reynolds RD, Lickteig JA, Deuster PA, Howard MP, Conway JM, Pietersma A, et al. Energy metabolism increases and regional body fat decreases while regional muscle mass is spared in humans climbing Mt. Everest. *J Nutr*. 1999;129:1307-14.

9. Bailey DM, Davies B, Milledge JS, Richards M, Williams SR, Jordinson M, et al. Elevated plasma cholecystokinin at high altitude: metabolic implications for the anorexia of acute mountain sickness. *High Alt Med Biol.* 2000;1:9-23.
10. Westerterp KR, Kayser B. Body mass regulation at altitude. *Eur J Gastroenterol Hepatol.* 2006;18(1):1-3.
11. Hamad N, Travis SP. Weight loss at high altitude: pathophysiology and practical implications. *Eur J Gastroenterol Hepatol.* 2006;18(1):5-10.
12. Roberts AC, Butterfield GE, Cymerman A, Reeves JT, Wolfel EE, Brooks GA, et al. Acclimatization to 4,300-m altitude decreases reliance on fat as a substrate. *J Appl Physiol.* 1996;81:1762-71.
13. McClelland GB, Hochachka PW & Weber JM. Carbohydrate utilization during exercise after high-altitude acclimation: a new perspective. *Proc Natl Acad Sci USA.* 1998;95:10288-93.
14. Viitala P, Newhouse IJ, Vitamin E supplementation, exercise and lipid peroxidation in human participants. *Eur J Appl Physiol.* 2004;93:108-15.
15. Schmidt MC, Askeew EW, Roberts DE, Prior RL, Ensign WY Jr, Hesslink RE Jr. Oxidative stress in humans training in a cold, moderate altitude environment and their response to a phytochemical antioxidant supplement. *Wilderness Environ Med.* 2002;13:94-105.
16. Streb P, Aubert S, Gout E, Bligny R. Reversibility of cold- and light-stress tolerance and accompanying changes of metabolite and antioxidant levels in the two high mountain plant species *Soldanella alpina* and *Ranunculus glacialis*. *J Exp Bot.* 2003;54:405-18.
17. Muzika RM, Guyette RP, Zielonka T, Liebhold AM. The influence of O₃, NO₂ and SO₂ on growth of *Picea abies* and *Fagus sylvatica* in the Carpathian Mountains. *Environ Pollut.* 2004;130:65-71.
18. Subudhi AW, Jacobs KA, Hagobian TA, Fattor JA, Fulco CS, Muza SR, et al. Antioxidant supplementation does not attenuate oxidative stress at high altitude. *Aviat Space Environ Med.* 2004;75:881-8.
19. Gámez A, Casas H, Panisello P, Viscor G, Pagés T, Carbonell T. Intermittent exposure to high altitude induces oxidative stress. *Proceedings of V World Congress of Mountain Medicine and High Altitude Physiology*; 2002 Apr 18-22; Barcelona, España: 2002. p. 18-27.
20. Bakonyi T, Radak Z. High altitude and free radicals. *J Sports Sci Med.* 2004;3:64-9.
21. Berglund B. High-altitude training. Aspects of haematological adaptation. *Sports Med.* 1992;14:289-303.
22. Heinicke K, Heinicke I, Schmidt W, Wolfarth B. A three-week traditional altitude training increases hemoglobin mass and red cell volume in elite biathlon athletes. *Int J Sports Med.* 2005;26(5):350-5.
23. Wehrlin JP, Zuest P, Hallén J, Marti B. Live high-train low for 24 days increases hemoglobin mass and red cell volume in elite endurance athletes. *J Appl Physiol.* 2006;100(6):1938-45. Epub 2006 Feb 23.
24. FAO/WHO Human energy requirements. Report of a Joint FAO/WHO/UNU Expert Consultation Rome, 2001.
25. Jiménez Cruz A, Cervera Ral P, Bacardi Gascón M. 2001 Novartis-Dietsource version 1.2. q0105071807.
26. Food Composition Table for Use in East Asia. Rome: Food Policy and Nutrition Division, Food and Agriculture Organization of the United Nations, 1972.
27. Westerterp KR. Limits to sustainable human metabolic rate. *J Exp Biol.* 2001;204(Pt 18):3183-7.
28. Dietary Reference Intakes for Energy, Carbohydrate, Fiber, Fat, Fatty Acids, Cholesterol, Protein, and Amino Acids (Macronutrients). National Academy of Sciences. Institute of Medicine. Food and Nutrition Board. 2005.
29. Altman DG, Bland JM. Statistics notes: the normal distribution. *BMJ.* 1995;310(6975):298.
30. Martínez González MA, Sánchez- Villegas A, Faulin-Fajardo J. *Bioestadística Amigable*. Madrid: Díaz de Santos; 2006.
31. Westerterp-Plantenga MS, Westerterp KR, Rubbens M, Verwegen CR, Ri-chelet JP, Gardette B. Appetite at "high altitude" [Operation Everest III (Comex-'97)]: a simulated ascent of Mount Everest. *J Appl Physiol.* 1999;87(1):391-9.
32. Brown MD, Dengel DR, Hogikyan RV, Supiano MA. Sympathetic activity and the heterogenous blood pressure response to exercise training in hypertensives. *J Appl Physiol.* 2002;92(4):1434-42.
33. Veitl V. Optimized nutrition for alpine athletes. *Wien Med Wochenschr.* 2000;150:191-4.
34. Askeew EW. Food for high-altitude expeditions: Pugh got it right in 1954 a commentary on the report by L.G.C.E. Pugh: "Himalayan rations with special reference to the 1953 expedition to Mount Everest". *Wilderness Environ Med.* 2004;15:121-4.
35. Bailey DM, Ainslie PN, Jackson SK, Richardson RS, Ghatei M. Evidence against redox regulation of energy homeostasis in humans at high altitude. *Clin Sci.* 2004;107:589-600.
36. Soares EA, Ishii M, Burini RC. Anthropometric and dietetic study of competitive swimmers from metropolitan areas of the southeastern region of Brazil. *Rev Saude Pub.* 1994;28:9-19.
37. Gleeson M, Bishop NC. Elite athlete immunology: importance of nutrition. *Int J Sports Med.* 2000;21:44-50.
38. Schena F, Guerrini F, Tregnaghi P, Kaiser B. Branched-chain amino acid supplementation during trekking at high altitude. The effects on loss of body mass, body composition, and muscle power. *Eur J Appl Physiol Occup Physiol.* 1992;65:394-8.
39. Fogelholm M. Dairy products, meat and sports performance. *Sports Med.* 2003;33:615-31.
40. Major GC, Doucet E. Energy intake during a typical Himalayan trek. *High Alt Med Biol.* 2004;5:355-63.
41. Paschoal VC, Amancio OM. Nutritional status of Brazilian elite swimmers. *Int J Sport Nutr Exerc Metab.* 2004;14:81-94.
42. Lukaski HC, Bolonchuk WW, Klevay LM, Milne DB, Sandstead HH. Interactions among dietary fat, mineral status, and performance of endurance athletes: a case study. *Nutr Exerc Metab.* 2002;12:381-3.
43. Nagyova A, Haban P, Klvanova J, Kadrabova J. Effects of dietary extra virgin olive oil on serum lipid resistance to oxidation and fatty acid composition in elderly lipidemic patients. *Bratisl Lek Listy.* 2003;104:218-21.
44. Vissers MN, Zock PL, Katan MB. Bioavailability and antioxidant effects of olive oil phenols in humans: a review. *Eur J Clin Nutr.* 2004;58:955-65.
45. Position of Dietitians of Canada, the American Dietetic Association, and the American College of Sports Medicine: Nutrition and Athletic Performance. (No authors listed). *Can J Diet Pract Res.* 2000;61:176-92.
46. Askeew EW. Work at high altitude and oxidative stress: antioxidant nutrients. *Toxicology.* 2002;180:107-19.
47. Nasolodin VV, Gladkikh IP, Meshcheriakov SI. Providing athletes with trace elements during intensive exercise. *Gig Sanit.* 2001;1:54-7.
48. Boning D, Cristancho E, Serrato M, Reyes O, Mora M Coy L, Rojas J. Hemoglobin mass and peak oxygen uptake in untrained and trained female altitude residents. *Int J Sports Med.* 2004;25:561-8.
49. Hinton PS, Sinclair LM. Iron supplementation maintains ventilatory threshold and improves energetic efficiency in iron-deficient nonanemic athletes. *Eur J Clin Nutr.* 2007;61(1):30-9. Epub 2006 Jul 12.
50. Bryant RJ, Ryder J, Martino P, Kim J, Craig BW. Effects of vitamin E and C supplementation either alone or in combination on exercise-induced lipid peroxidation in trained cyclists. *J Strength Cond Res.* 2003;17:792-800.
51. Thompson D, Williams C, McGregor SJ, Nicholas CW, McArdle F, Jackson MJ, et al. Prolonged vitamin C supplementation and recovery from demanding exercise. *Int J Sport Nutr Exerc Metab.* 2001;11:466-81.
52. Urso ML, Clarkson PM. Oxidative stress, exercise, and antioxidant supplementation. *Toxicology.* 2003;189:41-54.
53. Tur JA, Romaguera D, Pons A. Food consumption patterns in a Mediterranean region: does the Mediterranean diet still exist? *Ann Nutr Metab.* 2004;48:193-201.
54. Kayser B. Nutrition and energetics of exercise at altitude. Theory and possible practical implications. *Sports Med.* 1994;17(5):309-23.