



Junta de Andalucía  
Consejería de Educación y Deporte

# Revista Andaluza de Medicina del Deporte

<https://www.juntadeandalucia.es/deporte/ramd/>



Original

## Relationship between anaerobic work capacity and critical oxygenation in athletes



A. Vasquez-Bonilla<sup>a,\*</sup>, P. Tomas-Carus<sup>b,c</sup>, J. Brazo-Sayavera<sup>d</sup>, J. Malta<sup>b,c</sup>,  
H. Folgado<sup>b,c</sup>, G. Olcina<sup>a</sup>

<sup>a</sup> Research Group on Advances in Sport Training and Physical Conditioning, Faculty of Sport Science, University of Extremadura, Spain.

<sup>b</sup> Departamento de Desporto e Saúde, Escola de Ciências e Tecnologia, Universidade de Évora, Évora, Portugal.

<sup>c</sup> Comprehensive Health Research Centre (CHRC), University of Évora, Portugal.

<sup>d</sup> Universidad Pablo de Olavide, Sevilla, Spain.

ARTICLE INFORMATION: Received 17 September 2021, accepted 30 August 2022, online 2 September 2022

### ABSTRACT

**Objective:** Anaerobic work capacity (AWC) is understood as the maximum power that the athlete can withstand over time, conditioned by high intensity effort and it is important to interpret it for the performance improvement. In addition, the muscle oxygen saturation ( $SmO_2$ ) provides information on muscle metabolism and hemodynamics. Likewise, critical oxygenation (CO) is the highest metabolic rate that results in a fully oxidative energy supply that reaches a stable state at the substrate level. The main problem is that  $SmO_2$  generally offers a traditional laboratory interpretation without application in field tests. Therefore, the purpose of this study is to provide the use of CO as an indicator of AWC performance in high intensity exercise.

**Methods:** Twenty-two male rugby players participated. Peak torques during an isokinetic fatigue test and muscle oxygen consumption ( $mVO_2$ ) and  $SmO_2$  in the vastus lateralis were measured. A correlation and multiple regression analysis were applied to find an explanatory prediction model of the AWC.

**Results:** A greater  $SmO_2$  amplitude and CO would mean less anaerobic work ( $r = -0.58$  and  $r = -0.63$ ) and less force production. In addition, CO along with weight (kg) can explain the AWC by 64% during high intensity exercise.

**Conclusion:** The measurement of critical oxygenation is associated with the AWC, so should be considered a performance factor. These parameters could be included in NIRS sensors to evaluate muscle metabolism.

**Keywords:** Athletic performance; Skeletal muscle; Oxygen consumption; Anaerobic threshold; Energy metabolism; Regional blood flow.

## Relación entre la capacidad de trabajo anaeróbico y la oxigenación crítica en deportistas

### RESUMEN

**Objetivo:** La capacidad de trabajo anaeróbico (AWC) se entiende como la potencia máxima que el deportista puede soportar a lo largo del tiempo, condicionada por un esfuerzo de alta intensidad y es importante interpretarla para la mejora del rendimiento. Además, la saturación de oxígeno muscular ( $SmO_2$ ) proporciona información sobre el metabolismo muscular y la hemodinámica. Asimismo, la oxigenación crítica (OC) es la tasa metabólica más alta que da como resultado un suministro de energía completamente oxidativo que alcanza un estado estable a nivel de sustrato. El principal problema es que  $SmO_2$  generalmente ofrece una interpretación de laboratorio tradicional sin aplicación en pruebas de campo, por lo tanto, el propósito de este estudio es proporcionar el uso de OC como indicador del rendimiento de AWC en ejercicio de alta intensidad.

**Métodos:** Participaron 22 jugadores masculinos de rugby. Se midieron los picos máximos después de una prueba de fatiga isocinética y el consumo de oxígeno muscular ( $mVO_2$ ) y  $SmO_2$  en el músculo vasto lateral. Se aplicó un análisis de correlación y regresión múltiple para encontrar un modelo de predicción explicativo del AWC.

**Resultados:** Una mayor amplitud de  $SmO_2$  y OC supondría un menor trabajo anaeróbico ( $r = -0,58$  y  $r = -0,63$ ) y una menor producción de fuerza. Además, el CO junto con el peso (kg) pueden explicar el AWC en un 64 % durante el ejercicio de alta intensidad.

**Conclusión:** La medición de la oxigenación crítica está asociada a la AWC, por lo que debe considerarse un factor de rendimiento. Estos parámetros podrían incluirse en sensores NIRS para valorar el metabolismo muscular.

**Palabras claves:** Rendimiento atlético; Músculo esquelético; Consumo de oxígeno; Umbral anaeróbico; Metabolismo energético; Flujo sanguíneo regional.

\* Corresponding author.

E-mail-address: [alvasquezb@unex.es](mailto:alvasquezb@unex.es) (A. Vasquez-Bonilla).

<https://doi.org/10.33155/j.ramd.2022.09.001>

e-ISSN: 2172-5063/ © 2022 Consejería de Educación y Deporte de la Junta de Andalucía. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

## Relação entre capacidade de trabalho anaeróbico e oxigenação crítica em atletas

### RESUMO

**Objetivo:** A capacidade anaeróbica de trabalho (AWC) é entendida como a potência máxima que o atleta pode suportar ao longo do tempo, condicionada por um esforço de alta intensidade, sendo importante interpretá-la para melhorar o desempenho. Além disso, a saturação muscular de oxigênio ( $SmO_2$ ) fornece informações sobre o metabolismo muscular e a hemodinâmica. Da mesma forma, a oxigenação crítica (OC) é a taxa metabólica mais alta que resulta em um suprimento de energia totalmente oxidativo atingindo um estado estável no nível do substrato. O principal problema é que o  $SmO_2$  geralmente oferece uma interpretação laboratorial tradicional sem aplicação em testes de campo, portanto, o objetivo deste estudo é fornecer o uso do CO como indicador de desempenho de AWC em exercícios de alta intensidade.

**Métodos:** Participaram 22 jogadores de rugby do sexo masculino. Foram medidos os picos máximos após um teste de fadiga isocinética e o consumo de oxigênio muscular ( $mVO_2$ ) e  $SmO_2$  no músculo vasto lateral. Uma análise de correlação e regressão múltipla foi aplicada para encontrar um modelo explicativo de predição do AWC.

**Resultados:** Uma maior amplitude de  $SmO_2$  e CO implicaria em menor trabalho anaeróbico ( $r = -0,58$  e  $r = -0,63$ ) e menor produção de força. Além disso, o CO junto com o peso (kg) pode explicar a AWC em 64% durante o exercício de alta intensidade.

**Conclusão:** A medida oxigenação crítica prevê AWC, portanto, deve ser considerada um fator de desempenho. Esses parâmetros podem ser incluídos em sensores NIRS para a medição do metabolismo muscular.

**Palavras-chave:** Desempenho atlético; Músculo esquelético; Consumo de oxigênio; Limiar anaeróbico; Metabolismo energético; Fluxo sanguíneo regional.

### Introduction

The Anaerobic work capacity (AWC) is understood as the maximum power that the athlete can withstand over time, conditioned by high intensity effort.<sup>1</sup> Based on the maximum effort and duration of the exercise, the AWC is considered as an "oxygen independent" energy system where the phosphate and glycolysis pathway predominate, and to a lesser extent the oxidative phosphorylation.<sup>2,3</sup> The AWC performance is present in exercise modalities such as intermittent sprinting, maximum speed, repeated jumps and maximum strength exercises. In addition, AWC is related to the ability to oxygenate the muscles during high intensity exercise, due to the phosphocreatine (PCr) degradation process that is necessary to produce power.<sup>4</sup> So we have the possibility to predict the AWC over time through the amount of muscle oxygen consumption in athlete.<sup>5</sup>

Since the end of the 1980s to present, near infrared no-invasive (NIRS) technology has been used to assess the oxidative capacity of skeletal muscle through the consumption of muscle oxygen ( $mVO_2$ ) during exercise and in resting activities. The way it is generally measured is based on the differences in oxygenated hemoglobin ( $[O_2Hb]$  the one that is inside the muscle) and deoxygenated hemoglobin (Hbb) the one that the muscle uses).<sup>6</sup> However, the most commonly used NIRS-derived variable in field tests is muscle oxygen saturation ( $SmO_2$ ).<sup>7</sup>  $SmO_2$  reflects the dynamic balance between  $O_2$  supply and  $O_2$  consumption and is independent of the path of the near infrared photon in muscle tissue; its values range from 1% to 100%. There are several portable NIRS devices that use  $SmO_2$ . Studies have revealed that the lower the  $SmO_2$  values, greater the muscle oxygen extraction due to training load, but their interpretation is unknown in AWC activities. In this context  $SmO_2$  measurement could provide important information similar to  $mVO_2$ , observed through the difference between  $SmO_2$  at the exercise beginning divided by  $SmO_2$  at the exercise end ( $\Delta SmO_2$ ), this is known as the amplitude method.<sup>8</sup> It also possible to measure the time in which the  $SmO_2$  minimum values are reached as a performance and health factor.<sup>6</sup> Recently, the critical oxygenation (CO) model was published by Feldmann and Erlacher,<sup>9</sup> which represents "the greatest metabolic rate that results in wholly oxidative energy provision, and means that energy supply through substrate-level phosphorylation reaches a steady-state". The CO can be obtained through the  $SmO_2$  values at the beginning and the end exercise.

Studies carried out by Bosquet in 2015 and 2016 reveal that a specific test of 30 contractions (approximately 38 seconds) at the peak torque isokinetic could be interpreted as AWC values.<sup>10,11</sup> But the peak torque is an indirect calculation of the anaerobic metabolism in the absence of laboratory instruments to measure

energy pathways and oxygen consumption. So, an alternative to predict the AWC values could be the muscular oxygen consumption ( $mVO_2$ ) with a linear regression analysis from the beginning to the end of the exercise (point to point), this is a methodology known as "slope"<sup>8</sup> and is accurate, however it is difficult to take to daily training due to the lack of clarity to interpret the data in a field of play environment. However, the CO is a new model much easier to interpret in the field, but its relationship to the AWC is unknown. We hypothesize that CO has the ability to explain the AWC performance in athletes due to its close relationship with critical power measured in previous studies.<sup>9,12</sup> The aim of this study is report the relationship between CO and AWC during high intensity tests.

### Methods

#### Participants

A total of twenty-two male players (age  $22.5 \pm 4.6$  years, weight  $89.8 \pm 12.6$  kg, height  $176.4 \pm 7.8$  cm, sports participation  $9.1 \pm 3.6$  years) from the rugby Portuguese first division participated in the present study by signing a written statement of informed consent. To avoid any residual fatigue induced by recent training, participants were asked to refrain from strenuous exercise 48 hours before the tests. In addition, the research protocol was approved by the Scientific and Ethical Committee of the University of Extremadura with Number of registration: 131/2018 and it was in accordance with the principles of the Declaration of Helsinki.

#### Measures

- High-intensity isokinetic fatigue test (FAT)  
The test was performed on a Biodex System-3 Isokinetic Dynamometer (Biodex, System 3, NY, USA), which consisted of a 5-minute warm-up by pedaling at 100 watts with a cadence of around 100 revolutions per minute (rpm) on a cycloergometer (ergometrics 900, ergoline Germany). Thereafter, the participant was seated on the dynamometer seat, which was adjusted as previously described.<sup>13</sup> The range of motion was  $100^\circ$  ( $0^\circ$  corresponding to a full active extension). Afterwards, they performed 30 consecutive maximal reciprocal concentric contractions at an angular velocity of  $180^\circ/s$ . The work performed during the entire range of motion of each repetition was computed using the device's software and summed to obtain the FAT total work, peak moments (PM) torque was obtained for maximum and average, were taken in Newton-meter (N.M.).

- **Portable NIRS**  
SmO<sub>2</sub>, mVO<sub>2</sub> and CO values were measured using the portable NIRS sensor (Moxy, Fortiori Design LLC, Minnesota, USA). The device was placed in the vast lateral quadriceps between the greater trochanter and the lateral femoral epicondyle. For the analysis, data were recorded (visible only for researchers) using the raw SmO<sub>2</sub> and point to point Hbb and O<sub>2</sub>Hb signal was treated with a soft spline filter to reduce noise created by movement<sup>14</sup> Matlab® software (The MathWorks, Inc., Massachusetts, United States) and SmO<sub>2</sub> was observed in real time with ANT + technology (GoldenCheetah version 3.4, U.S.)
- **NIRS - Data Analysis**

The mVO<sub>2</sub> these values were compared with the values obtained from the linear regression of oxygenated hemoglobin (HbO) divided by deoxygenated hemoglobin (Hbb) that indicates the true mVO<sub>2</sub>. The SmO<sub>2</sub> slope was obtained from the difference between the SmO<sub>2</sub> value at the start of exercise and the muscle oxygen extraction (SmO<sub>2</sub> minimum value) during the FAT test (Figure 1). additionally, an expert assessed the CO based on the model proposed Feldmann and Erlacher,<sup>9</sup> finally, the times in sec of each SmO<sub>2</sub> point (a, b and c in Figure 1) were obtained as variables indicating a better oxidative performance.<sup>15</sup>


The formulas for each variable are presented below:

$$mVO_2 = HbO - Hbb \text{ constast}$$

$$SmO_2 \text{ slope} = SmO_2 \text{ Start} - SmO_2 \text{ minimum (from point (a) to point (b))}$$

$$CO \text{ model} = SmO_2 \text{ slope} - SmO_2 \text{ maximum (point (c))}$$


$$CO \text{ time (s)} = SmO_2 \text{ maximum time (s)} - SmO_2 \text{ minimum time (s)}$$



**Figure 1.** Dynamics of strength and SmO<sub>2</sub>% during a participant's evaluation in the dominant leg. (a)= Start of Exercise; (b)= Muscle oxygen extraction (Minimum SmO<sub>2</sub>); The SmO<sub>2</sub> Slope is the difference between point (a) to point (b). Intramuscular oxygen Partial pressure (PO<sub>2</sub>). (c)= Critical Oxygenation (Maximum SmO<sub>2</sub> after minimum SmO<sub>2</sub>).

Figure 2 shows how energy changes are obtained through the portable NIRS signal (MOXY), similar to the physiological mechanisms explained in the study by Conley et al.,<sup>16</sup> the review by Baker et al.<sup>17</sup> Sustainable AWC is determined at the muscular level, by the ability to maintain the ATP supply by decomposing PCr and by the products of glycolysis, which can inhibit the signal of oxidative phosphorylation, when exercise is very intense and therefore oxygen independent. This argument is based on magnetic resonance spectroscopy measurements of energy sources and sinks in vivo in human muscle and rattlesnake muscle during sustained contractions.<sup>16</sup> Likewise, the relationship between PO<sub>2</sub> and PCr can be explained by the balance maintained by creatine kinase, because a direct relationship was also found

between the relationship [ATPI/ADPI] with PCr.<sup>18</sup> Sympathetic vasoconstriction is attenuated by higher PO<sub>2</sub> and blood flow, which is related to a higher oxidative rate that reduces lactate production.<sup>19</sup>



**Figure 2.** Interaction of hemoglobin in the muscle during "Anaerobic Work Capacity" exercise and the change of the energy pathways.

**Statistical Analysis**

SPSS software (IBM® SPSS® Statistics 22) was used to perform the statistical analysis. Data are presented as means and standard deviation (SD). Multiple linear regressions with Pearson's coefficients were also used to establish the respective relationships between NIRS parameters and AWC. The following criteria were adopted for interpreting the magnitude of correlation (r) between test measures: 0.1, trivial; 0.1–0.3, small; 0.3–0.5, moderate; 0.5–0.7, large; 0.7–0.9, very-large; and 0.9–1.0, almost perfect. Regression analysis is a statistical technique for determining the relationship between a single dependent (criterion) variable and one or more independent (predictor) variables. The analysis yields a predicted value for the criterion resulting from a linear combination of the predictors. Statistical significance was set up with p value <0.05.

**Results**

Table 1 shows the mean results obtained during the FAT test. Table 2 as expected, there was a high correlation between mVO<sub>2</sub> and SmO<sub>2</sub> slope (r= 0.96), and CO correlation with mVO<sub>2</sub> (0.85) and SmO<sub>2</sub> slope (0.90). Furthermore, a greater amplitude of the SmO<sub>2</sub> slope would mean less anaerobic work (r = -0.58) and less force production. Also, a higher CO and CO time value is displayed with a lower AWC production (-0.63 and -0.76).

**Table 1.** Muscle oxygenation parameters obtained during the Anaerobic Work Capacity test.

ACW parameters	Mean ± SD
Peak Torque Max (N.m)	220 ± 30
Peak Torque Med (N.m)	170 ± 38
Total Work (Constant)	-3.01 ± 0.53
NIRS parameters	Mean ± SD
SmO <sub>2</sub> Start %	59.8 ± 7.4
SmO <sub>2</sub> minimum %	8.4 ± 14.1
SmO <sub>2</sub> medium %	21.1 ± 13.8
Critical Oxygenation %	20.8 ± 5.1
Critical Oxygenation % (seconds)	34 ± 8
SmO <sub>2</sub> Slope %	49.5 ± 14.9
mVO <sub>2</sub> (HbO/Hbb)	2.06 ± 0.53

Values are expressed as mean ± standard deviation.

Table 3 presents the multiple regression analysis models based on the SmO<sub>2</sub> dynamics. Moderate percentage correlations were observed in the equations to explain AWC by critical oxygenation. The use of time and weight are variables that explain the AWC by 64% and total work by 40%.

**Table 2.** Correlation between mVO<sub>2</sub>, SmO<sub>2</sub> slope, CO and AWC during high-intensity isokinetic exercise.

Variables	AWC	mVO <sub>2</sub>	SmO <sub>2</sub> Slope	CO	CO time
AWC	-	-	-	-	-
mVO <sub>2</sub>	-0.60 0.021*	-	-	-	-
SmO <sub>2</sub> slope	-0.58 0.037*	0.96 0.00**	-	-	-
CO	-0.63 0.016*	0.86 0.00**	0.90 0.00**	-	-
CO time	-0.76 0.00**	0.22 3.06	0.52 0.032*	0.54 0.029*	-

\*p value<0.05 and \*\*p value <0.01 statistically significant. Qualitative interpretation of the correlation magnitude (r) between test measures: 0.1, trivial; 0.1–0.3, small; 0.3–0.5, moderate; 0.5–0.7, large; 0.7–0.9, very-large; and 0.9–1.0, almost perfect.

Results are expressed in the procedure to calculate the AWC using the ΔSmO<sub>2</sub> data in athletes.

An example to evaluate AWC tests with the CO and CO time.

- $CO = SmO_2 \text{ Slope} - SmO_2 \text{ maximum during the test (1)}$
- $CO \text{ time} = SmO_2 \text{ maximum time} - SmO_2 \text{ minimum time (2)}$
- $Weight = \text{the subject's weight in kilograms}$
- $AWC = 99.34 + 0.38 * CO (SmO_2\%) - 2.40 * CO \text{ time} + 1.31 * weight (kg) (3)$
- $Total \text{ Work} = -2.07 + 0.03 * 1.02 * CO (SmO_2\%) - 0.04 * CO \text{ time} + 0.06 * weight (kg) (4)$

Example: a speed athlete weighting 64 kg runs a 400-meter test using a NIRS portable. The initial value is 65% (SmO<sub>2</sub> start) during the test and it is observed that the SmO<sub>2</sub> decreases to 26% (SmO<sub>2</sub> minimum) at the time of 26 seconds (SmO<sub>2</sub> minimum time). We would observe a SmO<sub>2</sub> Slope of 47%. Then a small increase and SmO<sub>2</sub> stabilization is observed at a value of 31% (SmO<sub>2</sub> maximum during the test). The test ended with a time of 48 seconds (SmO<sub>2</sub> maximum time).

From here we replace values with the formula:


$CO = 24\%$  of SmO<sub>2</sub>

$CO \text{ time} = 12 \text{ sec}$

$AWC = 99.34 + (0.38 * 24) - (2.40 * 12) + (1.31 * 64 \text{ kg})$

The AWC is 221 N.m

Answer: The athlete got an AWC of 221 N.m, presumably using the phosphocreatine energy system and glycolysis only in the first 36 sec. This procedure is using the CO model (Figure 3).



**Figure 3.** Critical oxygenation model to identify AWC through SmO<sub>2</sub> values

**Table 3.** Multiple regression analysis models for mVO<sub>2</sub> and AWC based on the measurement of SmO<sub>2</sub> Slope and CO with portable NIRS.

Step prediction models and statistical analysis	Independent variable	Dependent variable	Non-standardized coefficient B	Standardized coefficient B	Prediction Percentage	F Value	Sig.
First order equation Linear regression	SmO <sub>2</sub> Slope	mVO <sub>2</sub>	0.020 0.041	0.920	r= 0.92 r <sup>2</sup> = 85%	203.194	0.000**
First order equation Linear regression	SmO <sub>2</sub> Slope	AWC (N.m)	4.871 -0.044	-0.570	r= 0.57 r <sup>2</sup> = 32%	14.939	0.001*
First order equation Multiple linear regression	SmO <sub>2</sub> Slope Weight (kg)	AWC (N.m)	86.129 1.508 -0.243	0.504 -0.089	r= 0.60 r <sup>2</sup> = 36%	7.955	0.033*
First order equation Multiple linear regression	Time weight (kg) SmO <sub>2</sub> Slope	Total Work (N.m)	1.834 0.075 0.017 -0.041	0.220 0.163 -0.425	r= 0.54 r <sup>2</sup> = 29%	5.481	0.095
First order equation Multiple linear regression	Time weight (kg) SmO <sub>2</sub> Slope	AWC (N.m) when is "oxygen independent"	44.296 3.665 1.185 -0.562	0.377 0.396 -0.206	r= 0.65 r <sup>2</sup> = 43%	4.504	0.016*
First order equation Multiple linear regression	CO CO time Weight (kg)	Total Work (N.m)	-2.070 0.031 -0.045 0.057	0.490 -0.262 0.49	r= 0.63 r <sup>2</sup> = 40%	3.721	0.032*
First order equation Multiple linear regression	CO CO time Weight (kg)	AWC (N.m) when is "oxygen independent"	99.334 0.383 -2.394 1.312	0.250 -0.577 0.521	r= 0.80 r <sup>2</sup> = 64%	10.058	0.017*

\*p value<0.05 and \*\*p value <0.01 statistically significant. Correlation coefficient R<sup>2</sup> shows the explanatory model variance.

## Discussion

This study shows how anaerobic work capacity and CO level are highly related to the development of power during the isokinetic test, which we presume is extracted in greater quantity from the energy of PCr and anaerobic glycolysis. This is a mechanism that can be measured by sports scientists and coaches during high intensity events.

There are many studies that support our results, the study by Denis et al.,<sup>20</sup> found a decrease in SmO<sub>2</sub> at the beginning of the isokinetic dynamometer extension test and later an increase due to a compensatory vasodilator effect to counteract the decrease in SmO<sub>2</sub>. The explanation for this phenomenon is that at the beginning of exercise, an increase in intramuscular pressure occurs, restricts blood flow.<sup>21</sup> Vascular dilation and increased blood flow in the muscle is then promoted by increased production of metabolites (eg, K<sup>+</sup>, lactic acid). There is also an increase in muscle oxygenation detected by NIRS, which reflects the volume of blood in small vessels, including arterioles, capillaries and venules.<sup>22</sup> The temperature of the muscles and skin may have increased after the start of exercise, which also leads to dilation of the microvasculature.<sup>22</sup> This allows the acute positive regulation of vasodilator factors such as adenosine and nitric oxide (NO),<sup>23</sup> which respond to low local PO<sub>2</sub>.<sup>24,25</sup> Low PO<sub>2</sub> values strongly affect the O<sub>2</sub>-ATP reaction rate, but does not significantly compromise the use of intramuscular oxygen<sup>26</sup> and therefore does not substantially limit the consumption of oxygen. Rather deoxygenation under anaerobic conditions occurs by the transport of the myoglobin molecule, which also supports the diffusion-facilitating role that Mb can play in O<sub>2</sub> transport because Mb needs to be desaturated to facilitate the movement of O<sub>2</sub> from the blood to the cell.<sup>26</sup>

A lower intramuscular PO<sub>2</sub> also turns off phosphorus and affects the NIRS signal<sup>27</sup> which hinders myosin-actin bridge formation, and initiates the loss of contractile efficiency and force generation. Gomez-Carmona et al.,<sup>28</sup> have determined the oxygen differences as force production and found that, over time, they decrease. Generally, these differences in SmO<sub>2</sub> Slope are accompanied by the use of glycolysis to generate ATP, because arterial epinephrine levels stimulate glycolysis and are closely related to skeletal muscle lactate output.<sup>18,29</sup> Likewise, the Feldmann (2021) study, which used SmO<sub>2</sub> in a 3 min "all out" test, showed progress in finding the critical power, where a phenomenon similar to our study was observed: muscle oxygen extraction and CO, referring to the increased metabolic rate resulting in a fully oxidative energy supply.<sup>9</sup> It is the same model used in this study.

Likewise, energy metabolism can be explained by theoretical studies such as that by Gatin et al.<sup>30</sup> and Barclay<sup>31</sup> who argued that such a short test of anaerobic work reflects a decrease in performance, possibly due to the change in the metabolic pathway within the muscle. Similarly, other studies have indicated that a greater contribution of the energy-oxidative phosphorylation pathways is connected with an hyperaemia increase and blood flow, which causes the muscle to have a vasodilator effect and increases SmO<sub>2</sub>.<sup>5,32</sup> In addition, during this metabolic process, the difference in the interindividual response of the subjects depends on the capacity of the cardiovascular system and the haemodynamics of the capillary beds to maintain the supply of oxygen in the muscle. This is observable in the almost complete desaturation in the anaerobic peak test which can occur due to relatively restricted blood flow.<sup>4,27</sup> Then, increased blood flow is a mechanism that attenuates sympathetic vasoconstriction in active muscles by metabolic events in contracting skeletal muscle, in part by the activation of ATP-sensitive potassium (KATP) channels. Sympathetic vasoconstriction is mediated by the endogenous vasodilator NO, which is necessary to optimize muscle O<sub>2</sub> perfusion.<sup>33,34</sup> It is important to clarify that the greater activation capacity of type II fibres to extract oxygen through the glycolytic

pathway is necessary to achieve better performance in high intensity areas and maintain a greater force and power production, because type II fibres need less oxygen to function.<sup>32</sup>

Many studies have observed a non-linear (hyperbolic) relationship of muscle oxygenation in high-intensity zones.<sup>5,35</sup> PO<sub>2</sub> is reduced and then remains constant despite a large continuous increase in work rate. Intracellular oxygenation does not, however, fall linearly with increasing metabolic rate, which indicates oxygen availability from submaximal exercise to single-leg knee extensor maximum.<sup>29</sup> Although the findings are straightforward, there are many far-reaching implications from these data. Changes in oxygenation levels over short periods have not been measurable with portable NIRS, and sometimes not with expensive instruments and laboratory tests. However, we can provide an explanation of this important phenomenon in measurements of oxidative metabolism at the local level, which could lead to better athletic performance. This may be a possible explanation for the lack of linearity. An important point is that improved muscle oxygenation during exercise leads to improved K<sup>+</sup> regulation in humans.<sup>36</sup>

Regarding mVO<sub>2</sub> and SmO<sub>2</sub>, during an exclusively anaerobic work test (sprint, jump, etc.), they report important information for the AWC assessment, however the objective of this study is to offer as an alternative analysis the "method of amplitude" which consists of taking the differences in the slope of SmO<sub>2</sub> up to the point at which it reaches the plateau<sup>8</sup> that is, at best power production. Also provide sufficient information on the CO model recently published by Feldmann<sup>9</sup> (Figure 3).

It is worth noting that our study is an explanatory model with multiple regressions; to date there are no studies that only compare anaerobic work with SmO<sub>2</sub>. In our study, we presumed that critical oxygenation could predict anaerobic work due to the similarity of the SmO<sub>2</sub> slopes and mVO<sub>2</sub> as variables identified in the analysis of muscle oxygen extraction;<sup>37</sup> however, that calculation is better when only the CO is used (Figure 3).

The main limitation of this study is that it does not measure the contribution of each energy pathway, as in the studies of Milioni et al.,<sup>38</sup> where a gas analyzer is needed, in addition to measurements of vascular dilation and muscle blood flow. However, we can overlap these data due to the high correlation of SmO<sub>2</sub> with VO<sub>2</sub> and blood flow.<sup>39</sup> This study proposes an advance in the physiology of muscle oxygenation that is applicable in a practical way by coaches in the field and by researchers to determine the athletes performance and to plan training.

Summary, the critical oxygenation is associated with AWC test and is a peripheral performance factor. In addition, this information could be used as a new portable NIRS methodology in sports science studies and by coaches assessing muscle metabolism in maximum speed and power tests, sprint trials, jumps and intermittent work intervals to scientifically support the sports training. Processed data can be exported to NIRS instruments or integrated into smartphone apps and software. However, this information must be handled with care by scientists, as it must be tested and supported with other anaerobic tests to improve the AWC calculation.

**Authorship.** All the authors have intellectually contributed to the development of the study, assume responsibility for its content and also agree with the definitive version of the article. **Conflicts of interest.** The authors have no conflicts of interest to declare. **Funding.** This study has been supported by the Government of Extremadura with funding from the European Regional Development Fund under grant (Ref: GR18003), QUERCUS + 2018/19 (Erasmus + Internships). **Provenance and peer review.** Not commissioned; externally peer reviewed. **Ethical Responsibilities.** *Protection of individuals and animals:* The authors declare that the conducted procedures met the ethical standards of the responsible committee on human experimentation of the World Medical Association and the Declaration of Helsinki. *Confidentiality:* The authors are responsible for following the protocols established by their respective healthcare centers for accessing data from medical records for performing this type of publication in order to conduct research/dissemination for the community. *Privacy:* The authors declare no patient data appear in this article.

## References

- Green S. [Measurement of anaerobic work capacities in humans.](#) *Sports Med.* 1995;19(1):32-42.
- Chamari K, Padulo J. ['Aerobic' and 'Anaerobic' terms used in exercise physiology: a critical terminology reflection.](#) *Sports Med Open.* 2015;1(1):9.
- Spencer MR, Gastin PB. [Energy system contribution during 200- to 1500-m running in highly trained athletes.](#) *Med Sci Sports Exerc.* 2001;33(1):157-62.
- Nioka S, Moser D, Lech G, Evangelisti M, Verde T, Chance B, et al. [Muscle deoxygenation in aerobic and anaerobic exercise.](#) *Adv Exp Med Biol.* 1998;454:63-70.
- Vasquez-Bonilla AA, Camacho-Cardeñosa A, Timón R, Martínez-Guardado I, Camacho-Cardeñosa M, Olcina G. [Muscle oxygen desaturation and re-saturation capacity limits in repeated sprint ability performance in women soccer players: a new physiological interpretation.](#) *Int J Environ Res Public Health.* 2021; 18(7):3484.
- Hamaoka T, McCully KK, Quaresima V, Yamamoto K, Chance B. [Near-infrared spectroscopy/imaging for monitoring muscle oxygenation and oxidative metabolism in healthy and diseased humans.](#) *J Biomed Opt.* 2007;12(6):062105.
- Celie B, Boone J, Van Coster R, Bourgois J. [Reliability of near infrared spectroscopy \(NIRS\) for measuring forearm oxygenation during incremental handgrip exercise.](#) *Eur J Appl Physiol.* 2012;112(6):2369-74.
- Agbangla NF, Audiffren M, Albinet CT. [Assessing muscular oxygenation during incremental exercise using near-infrared spectroscopy: comparison of three different methods.](#) *Physiol Res.* 2017;66(6):979-985.
- Feldmann A, Erlacher D. [Critical oxygenation: Can muscle oxygenation inform us about critical power?](#) *Med Hypotheses.* 2021;150:110575.
- Bosquet L, Gouadec K, Berryman N, Duclos C, Gremeaux V, Croisier JL. [The total work measured during a high intensity isokinetic fatigue test is associated with anaerobic work capacity.](#) *J Sports Sci Med.* 2016;15(1):126-30.
- Bosquet L, Gouadec K, Berryman N, Duclos C, Gremeaux V, Croisier JL. [Physiological interpretation of the slope during an isokinetic fatigue test.](#) *Int J Sports Med.* 2015;36(8):680-3.
- Kirby BS, Clark DA, Bradley EM, Wilkins BW. [The balance of muscle oxygen supply and demand reveals critical metabolic rate and predicts time to exhaustion.](#) *J Appl Physiol (1985).* 2021;130(6):1915-1927.
- Bosquet L, Maquet D, Forthomme B, Nowak N, Lehance C, Croisier JL. [Effect of the lengthening of the protocol on the reliability of muscle fatigue indicators.](#) *Int J Sports Med.* 2010 Feb;31(2):82-8.
- Rodriguez RF, Townsend NE, Aughey RJ, Billaut F. [Influence of averaging method on muscle deoxygenation interpretation during repeated-sprint exercise.](#) *Scand J Med Sci Sports.* 2018;28(11):2263-2271.
- Perrey S, Ferrari M. [Muscle oximetry in sports science: a systematic review.](#) *Sports Med.* 2018;48(3):597-616.
- Conley KE, Kemper WE, Crowther GJ. [Limits to sustainable muscle performance: interaction between glycolysis and oxidative phosphorylation.](#) *J Exp Biol.* 2001;204(Pt 18):3189-94.
- Baker JS, McCormick MC, Robergs RA. [Interaction among Skeletal Muscle Metabolic Energy Systems during Intense Exercise.](#) *J Nutr Metab.* 2010;2010:905612.
- Bylund-Fellenius AC, Walker PM, Elander A, Holm S, Holm J, Scherstén T. [Energy metabolism in relation to oxygen partial pressure in human skeletal muscle during exercise.](#) *Biochem J.* 1981;200(2):247-55.
- Rosenmeier JB, Hansen J, González-Alonso J. [Circulating ATP-induced vasodilatation overrides sympathetic vasoconstrictor activity in human skeletal muscle.](#) *J Physiol.* 2004;558(Pt 1):351-65.
- Denis R, Bringard A, Perrey S. [Vastus lateralis oxygenation dynamics during maximal fatiguing concentric and eccentric isokinetic muscle actions.](#) *J Electromyogr Kinesiol.* 2011;21(2):276-82.
- Crenshaw AG, Bronce L, Krag I, Jensen BR. [Oxygenation and EMG in the proximal and distal vastus lateralis during submaximal isometric knee extension.](#) *J Sports Sci.* 2010;28(10):1057-64.
- Watanabe S, Ishii C, Takeyasu N, Ajisaka R, Nishina H, Morimoto T, et al. [Assessing muscle vasodilation using near-infrared spectroscopy in cardiac patients.](#) *Circ J.* 2005;69(7):802-14.
- Kimura M, Ueda K, Goto C, Jitsuiki D, Nishioka K, Umemura T, et al. [Repetition of ischemic preconditioning augments endothelium-dependent vasodilation in humans: role of endothelium-derived nitric oxide and endothelial progenitor cells.](#) *Arterioscler Thromb Vasc Biol.* 2007 Jun;27(6):1403-10.
- Takakura H, Masuda K, Hashimoto T, Iwase S, Jue T. [Quantification of myoglobin deoxygenation and intracellular partial pressure of O<sub>2</sub> during muscle contraction during haemoglobin-free medium perfusion.](#) *Exp Physiol.* 2010;95(5):630-40.
- Richardson RS, Newcomer SC, Noyszewski EA. [Skeletal muscle intracellular PO<sub>2</sub> assessed by myoglobin desaturation: response to graded exercise.](#) *J Appl Physiol (1985).* 2001;91(6):2679-85.
- Groebe K, Thews G. [Calculated intra- and extracellular PO<sub>2</sub> gradients in heavily working red muscle.](#) *Am J Physiol.* 1990;259(1 Pt 2):H84-92.
- Koga S, Kano Y, Barstow TJ, Ferreira LF, Ohmae E, Sudo M, Poole DC. [Kinetics of muscle deoxygenation and microvascular PO<sub>2</sub> during contractions in rat: comparison of optical spectroscopy and phosphorescence-quenching techniques.](#) *J Appl Physiol (1985).* 2012 Jan;112(1):26-32.
- Gómez-Carmona CD, Bastida-Castillo A, Rojas-Valverde D, de la Cruz Sánchez E, García-Rubio J, Ibáñez SJ, et al. [Lower-limb Dynamics of Muscle Oxygen Saturation During the Back-squat Exercise: Effects of Training Load and Effort Level.](#) *J Strength Cond Res.* 2020 May;34(5):1227-1236.
- Richardson RS. [Oxygen transport: air to muscle cell.](#) *Med Sci Sports Exerc.* 1998 Jan;30(1):53-9.
- Gastin PB. [Energy system interaction and relative contribution during maximal exercise.](#) *Sports Med.* 2001;31(10):725-41.
- Barclay CJ. [Energy demand and supply in human skeletal muscle.](#) *J Muscle Res Cell Motil.* 2017;38(2):143-155.
- Inglis EC, Iannetta D, Murias JM. [The plateau in the NIRS-derived \[HHb\] signal near the end of a ramp incremental test does not indicate the upper limit of O<sub>2</sub> extraction in the vastus lateralis.](#) *Am J Physiol Regul Integr Comp Physiol.*

- [2017;313\(6\):R723-R729.](#)
33. [Murias JM, Spencer MD, Keir DA, Paterson DH. Systemic and vastus lateralis muscle blood flow and O2 extraction during ramp incremental cycle exercise. Am J Physiol Regul Integr Comp Physiol. 2013;304\(9\):R720-5.](#)
  34. [Thomas GD, Victor RG. Nitric oxide mediates contraction-induced attenuation of sympathetic vasoconstriction in rat skeletal muscle. J Physiol. 1998;506:817-26.](#)
  35. [Calbet JA, Gonzalez-Alonso J, Helge JW, Søndergaard H, Munch-Andersen T, Boushel R, et al. Cardiac output and leg and arm blood flow during incremental exercise to exhaustion on the cycle ergometer. J Appl Physiol \(1985\). 2007;103\(3\):969-78.](#)
  36. [Christiansen D. Molecular stressors underlying exercise training-induced improvements in K<sup>+</sup> regulation during exercise and Na<sup>+</sup>, K<sup>+</sup>-ATPase adaptation in human skeletal muscle. Acta Physiol. 2019;225\(3\): e13196.](#)
  37. [Paquette M, Bieuzen F, Billaut F. Muscle Oxygenation Rather Than VO2max as a Strong Predictor of Performance in Sprint Canoe-Kayak. Int J Sports Physiol Perform. 2018 Nov 19:1-9.](#)
  38. [Milioni F, Zagatto AM, Barbieri RA, Andrade VL, Dos Santos JW, Gobatto CA, da Silva AS, Santiago PR, Papoti M. Energy Systems Contribution in the Running-based Anaerobic Sprint Test. Int J Sports Med. 2017 Mar;38\(3\):226-232.](#)
  39. [Wolf M, Ferrari M, Quaresima V. Progress of near-infrared spectroscopy and topography for brain and muscle clinical applications. J Biomed Opt. 2007 Nov-Dec;12\(6\):062104.](#)